

4.1 Introduction

When Mozart performed his opera Die Entführung aus dem Serail, the Emperor Joseph II responded wryly, 'Too many notes, Mozart!'



In this chapter we meet a different problem: too many numbers!

We have met *discrete* random variables, for which we can *list all the values* Contable infinite and their probabilities, even if the list is infinite:

e.g. for $X \sim \text{Geometric}(p)$,

But suppose that X takes values in a *continuous set*, e.g. $[0, \infty)$ or (0, 1)

We can't even begin to list all the values that X can take. For example, how would you list all the numbers in the interval [0, 1]?

• the smallest number is 0, but what is the next smallest? 0.01? 0.0001? 0.0000000001? We just end up talking *nonsense*.

In fact, there are so many numbers in any continuous set that each of them must have probability 0.

If there was a probability > 0 for all the numbers in a continuous set, however 'small', there simply wouldn't be enough probability to go round.

A continuous random variable takes values in a continuous interval (a, b). It describes a continuously varying quantity such as time or height. When X is continuous, $\mathbb{P}(X = x) = 0$ for ALL x. The probability function is meaningless. Although we cannot assign a probability to any *value* of X, we *are* able to assign probabilities to *intervals*: eg. $\mathbb{P}(X = 1) = 0$, but $\mathbb{P}(0.999 \le X \le 1.001)$ can be > 0. interval probabilit

This means we should use the distribution function, $F_X(x) = \mathbb{P}(X \leq x)$.

$$\mathbb{E}\left(\frac{\beta}{\beta}\right) = \mathbb{E}\left(\frac{Y_{1} + Y_{2} + Y_{3}}{6}\right) \geq E\left(\frac{\beta}{2}\right) \leq E\left(\frac{\beta}{2}$$

 $Y_i \sim Poisson(\beta x_i)$



Discrete r.v.s. Fx is a STEP

Defn of a continuous r.v. Continuous V.V.S,

probability accumulates

continuously Horoughout range of X.

P(X=1)

ÌP(X=e)

 $F_X(x)$

Ω

The cumulative distribution function, $F_X(x)$

Recall that for *discrete* random variables:

- $F_X(x) = \mathbb{P}(X < x);$
- $F_X(x)$ is a step function: probability accumulates in discrete steps;

•
$$\mathbb{P}(a < X \leq b) = \mathbb{P}(X \in (a, b]) = F(b) - F(a)$$

For a *continuous random variable:* $F_X(x)$ 1

- $F_X(x) = \mathbb{P}(X < x)$:
- $F_X(x)$ is a <u>continuous function</u>: probability accumulates continuously;
- As before, $\mathbb{P}(a < X \leq b) = \mathbb{P}(X \in (a, b]) = F(b) F(a)$.

However, for a *continuous* random variable,

$$\mathbb{P}(X=a)=0.$$

So it makes no difference whether we say $\mathbb{P}(a < X \leq b)$ or $\mathbb{P}(a \leq X \leq b)$.



Endpoints are very important for discrete r.v.s.



The probability density function 4.2

Although the cumulative distribution function gives us an interval-based tool for dealing with continuous random variables, it is not very good at telling us what the distribution *looks like*. $f_{\mathbf{x}}(\mathbf{x})$

For this we use a different tool called the *probability density function*.

The probability density function (p.d.f.) is the best way to describe and recognise a continuous random variable. We use it all the time to calculate probabilities and to gain an intuitive feel for the shape and nature of the distribution. Using the p.d.f. is like recognising your friends by their faces. You can chat on the phone, write emails or send txts to each other all day, but you never really know a person until you've seen their face.

Just like a cell-phone for keeping in touch, the cumulative distribution function is a tool for facilitating our interactions with the continuous random variable. However, we never really understand the random variable until we've seen its 'face' — the probability density function. Surprisingly, it is quite difficult to describe exactly what the probability density function is. In this section we take some time to motivate and describe this fundamental idea.

All-time top-ten 100m sprint times

The histogram below shows the best 10 sprint times from the 168 all-time top male 100m sprinters. There are 1680 times in total, representing the top 10 times up to 2002 from each of the 168 sprinters. Out of interest, here are the summary statistics:







We could plot this histogram using different time intervals:



We see that each histogram has broadly the same shape, although the heights of the bars and the interval widths are different.

The histograms tell us the most intuitive thing we wish to know about the distribution: its *shape*:

- the *most probable* times are *close to 10.2 seconds*;
- the distribution of times has a *long left tail (left skew)*;
- times below 10.0s and above 10.3 seconds have *low probability*.

126



We could fit a curve over any of these histograms to show the desired shape, but the problem is that *the histograms are not standardized*:

• every time we change the interval width, *the heights of the bars change*.

How can we derive a curve or function that captures the common shape of the histograms, but keeps a constant height? What should that height be?

The standardized histogram

We now focus on an *idealized* (smooth) version of the sprint times distribution, rather than using the exact 1680 sprint times observed.

We are aiming to derive a curve, or function, that captures the shape of the histograms, but will keep the same height for any choice of histogram bar width.

<u>First idea:</u> plot the probabilities instead of the frequencies.

The height of each histogram bar now represents the probability of getting an observation in that bar.



This doesn't work, because the height (probability) still depends upon the bar width. Wider bars have higher probabilities.



This seems to be exactly what we need! *The same curve fits nicely over all the histograms and keeps the same height regardless of the bar width.*

These histograms are called *standardized histograms*.

The nice-fitting curve is the probability density function. $\int_{\mathcal{T}} (\mathbf{x})$.

But... what is it?!

The probability density function $f_{\mathbf{x}}(\mathbf{x}) = \mathbf{P}(\mathbf{x}=\mathbf{x})$. We have seen that there is a single curve that fits nicely over any standardized histogram from a given distribution. This curve is called the *probability density function* (p.d.f.). We will write the p.d.f. of a continuous random variable X as $PDF = f_{\mathbf{x}}(\mathbf{x})$. The p.d.f. $f_{\mathbf{x}}(\mathbf{x})$ is NOT the probability of $\mathbf{x}=\mathbf{x}$. For example, in the spint distribution we can have $f_{\mathbf{x}}(\mathbf{x}) = 4$, so it dearly can Not be a probability. Anyway, $P(\mathbf{x}=\mathbf{x})=0$ for all \mathbf{x} when X. However, as the histogram bars of the standardized histogram get narrower, *on the standardized histogram bars hight as the bar width approadus* What is the height of the standardized histogram bar? For an interval from \mathbf{x} to $\mathbf{x} + \mathbf{t}$, the standardized histogram plots the probability

For an interval from x to x+t, the standardized histogram plots the probability of an observation falling between x and x+t, divided by the width of the interval, t.

Thus the height of the standardized histogram bar over the interval from x to x + t is:

$$\frac{\text{probability}}{\text{intorval width}} = \frac{P(x \leq X \leq x+t)}{t} = \frac{F_{X}(x+t) - F_{X}(x)}{t}$$
Where $F_{X}(x)$ is the C.D.F.

Now consider the limit as the histogram bar width (t) goes to 0:

this limit is DEFINED TO BE the P.D.F. at x, fr (x):

$$f_{X}(x) = f_{im} \begin{cases} \frac{F_{X}(x+t) - F_{X}(x)}{t} \\ \frac{F_{X}(x+t) - F_{X}(x)}{t} \end{cases} \text{ by defn.}$$

This expression should look familiar: it is the derivative of $F_{X}(x)$: $F_{X}'(x)$.



The probability density function (p.d.f.) is therefore

 $f_{x}(x) = F_{x}'(x) = \frac{dF_{x}(x)}{dx}.$ It is a single, unchanging curve that describes the SHAPE of a histogram of observations drawn from the distr of X. Formal definition of the probability density function

Definition: Let X be a continuous random variable with distribution function $F_X(x)$. The **probability density function (p.d.f.)** of X is defined as

$$f_{x}(x) = \frac{dF_{x}(x)}{dx} = F_{x}'(x).$$

It gives:

- the RATE at which probability is accumulating at point x;
- · the SHAPE of the distribution of X.

Using the probability density function to calculate probabilities

As well as showing us the shape of the distribution of X, the probability density function has another major use:

•
$$f_{x}(x)$$
 provides probability calculations by integration.
Suppose we want to calculate $P(a \le X \le b)$.
We already know that:
 $P(a \le X \le b) = F_{x}(b) - F_{x}(a)$.
But we also know that:
 $\frac{dF_{x}}{dx} = f_{x}(x)$
So $F_{x}(x) = \int F_{x}(x) dx$ (without constants)
In fact: $F_{x}(b) - F_{x}(a) = \int_{a}^{b} f_{x}(x) dx$



131

This is a very important result:

Let X be a continuous random variable with probability density function $f_X(x)$. Then

$$\mathbb{P}(a \leq X \leq b) = \int_{a}^{b} \mathcal{F}_{x}(x) dx \cdot \frac{KNOU}{TH}.$$



The total area under the p.d.f. curve is:

$$f_{ofalorea} = \int_{-\infty}^{\infty} f_{x}(x) dx = F_{x}(\infty) - F_{x}(-\infty)$$

= 1 - 0 = 1.

This says that the total area under the p.d.f. curve is equal to the total probability that X takes a value between $-\infty$ and $+\infty$, which is



$$f_{X}(n) = \frac{d}{dn} f_{X}(n) \Longrightarrow f_{X}(n) = \int_{-\infty}^{x} f_{X}(n) dn \xrightarrow{\text{rev zealand}}_{\text{rev zealand}} \text{The UNIVERSITY}_{\text{rev zealand}}$$
Using the p.d.f. to calculate the distribution function, $F_{X}(x)$

Suppose we know the probability density function, $f_X(x)$, and wish to calculate the distribution function, $F_X(x)$. We use the following formula:

Comulative
distribution function,
$$f_{x}(x) = \int_{-\infty}^{x} f_{x}(u) du$$

Proof:

$$\int_{-\infty}^{x} f_{x}(u) du = F_{x}(x) - F_{x}(-\infty) = F_{x}(x) - 0 = F_{x}(x).$$

$$\int_{-\infty}^{x} f_{x}(u) du = F_{x}(u) \int_{-\infty}^{x} f_{x}(u) du$$
Using the domain unrights as

Using the dummy variable, u:



How can x range from $-\infty$ to x?!



These graphs show $F_X(x)$ and $f_X(x)$ from the men's 100m sprint times (X is a random top ten 100m sprint time).



Using the p.d.f., $f_X(x)$, we can see that it is about $|0 \cdot |$ to $|0 \cdot 2$ seconds.

Using the c.d.f., $F_X(x)$, we would have to inspect the part of the curve with the steepest gradient: very difficult to see.





 $-\frac{k}{2}\left(e^{-\omega}-e^{\omega}\right)$ 134=0 2. $\begin{cases} 2e^{-2x} \text{ for } 0 < x < \infty \\ 0 \end{cases}$ So $f_{x}(x)$ = 0 $-\infty < \chi \leq 0$. ii) $P(1 \le X \le 3) = \int_{1}^{3} F_{X}(n) dx$ $= \int_{1}^{3} 2e^{-2\pi} d\pi$ $= \left[-e^{-2x} \right]_{1}^{3}$ $=\frac{-2}{2}e^{-2*3}-e^{-2*1}$ 0.132 iii) $F_{x}(x) = \int_{x}^{\infty} f_{x}(u) du$ $f_{X_{\downarrow}}(n)$ 20 $= \int_{-\infty}^{\infty} 0 \, du + \int_{0}^{\infty} 2e^{-2u} \, du \quad \text{for } x > 0$ $0 du + \int_{0}^{2} 2e^{-2u} du = 0$ $+ \left[-e^{-2u} \right]_{0}^{2} \qquad So \quad \text{final answer:} \\ F_{x}(x) = \int_{0}^{1-e^{-2u}} for \ n > 0$ $= \int_{0}^{2-o} 2e^{-2u} \int_{0}^{2} for \ n > 0$ $= \int_{0}^{2-o} 2e^{-2u} \int_{0}^{2} for \ n > 0$ $- \xi e^{-2x} - e^{-0}$ 5 e-2x - 13 1- e-2x. for x>0 F_{\star} (x)





 $F_{\chi}(\chi) = \begin{cases} 0 & \text{for } \chi \leq 0 \\ \text{Llah2} & \text{for } 0 < \chi < 6 \\ 1 & \text{for } \chi \geq 6 \end{cases}$



4.3 The Exponential distribution



When will the next volcano erupt in Auckland? We never quite answered this question in Chapter 3. The Poisson distribution was used to count the hunder

of ruptions that will occur in a fixed space of time. We have not said how long we have to wait for the next volcano: this is a continuous random variable (measure of time).

Auckland Volcanoes

About 50 volcanic eruptions have occurred in Auckland over the last 100,000 years or so. The first two eruptions occurred in the Auckland Domain and Albert Park — right underneath us! The most recent, and biggest, eruption was Rangitoto, about 600 years ago. There have been about 20 eruptions in the last 20,000 years, which has led the Auckland Council to assess current volcanic risk by assuming that volcanic eruptions in Auckland follow a Poisson process with rate $\lambda = \frac{1}{1000}$ volcanoes per year. For background information, see: www.aucklandcouncil.govt.nz and search for 'volcanic hazard'.

Distribution of the waiting time in the Poisson process

The length of time between events in the Poisson process is called the Waiting time.

To find the distribution of a continuous random variable, we often work with the CDF, $F_{x}(x)$.

This is because $F_X(x) = \mathcal{P}(X \leq \varkappa)$ gives us a *probability*, unlike the p.d.f. $f_X(x)$. We are comfortable with handling and manipulating probabilities.

Suppose that $\{N_t : t > 0\}$ forms a Poisson process with rate $\lambda = \frac{1}{1000}$. N_t is the number of emptions to occur by time t, Starting now.

We know that $N_t \sim Poisson (\lambda t)$

So

$$\mathbb{P}\left(\mathsf{N}_{\mathsf{t}}=\mathsf{n}\right)=\frac{\left(\lambda\mathsf{t}\right)^{\mathsf{n}}}{\mathsf{n}!}e^{-\lambda\mathsf{t}}$$



Let X be a continuous random variable giving the years waited before the next supprise, starting now. We will derive an expression for $F_X(x)$.

(i) When
$$x < 0$$
:

$$F_{X}(x) = P(X \le x)$$

$$= P(less than 0 time before rext emplies)$$
(ii) When $x > 0$:
(i) When $x > 0$:



because of the exponential formula for $F_X(x)$.

Example: What is the probability that there will be a volcanic eruption in Auck-#Eruptions ~ Poisson Process, Z= 1000 land within the next 50 years? =) Time waited ~ Exponential $(\lambda = \frac{1}{1000})$ (in years). years. Let time waited, $X \sim Exponential \left(\frac{1}{1000}\right)$ We need $P(X \le 50) = F_x(50) = 1 - e^{-\frac{1}{1000} * 50} =$

There is about a 5% change that there will be a volcanic eruption in Auckland over the next 50 years. This is the figure given by the Auckland Council at the above web link.







Memorylessness

We have said that the waiting time of the Poisson process can be defined *either* as the time from the start to the first event, *or* the time from now until the next event, *or* the time between any two events.



All of these quantities have the same distribution: $X \sim \text{Exponential}(\lambda)$.

The derivation of the Exponential distribution was valid for all of them, because events occur at a constant average rate in the Poisson process.

This property of the Exponential distribution is called *memorylessness*:



Rangitoto erupted have counted for nothing.

The chance that we still have 1000 years to wait for the next eruption is the same today as it was 600 years ago when Rangitoto erupted.

Memorylessness applies to *any* Poisson process. It is not always a desirable property: you don't want a memoryless waiting time for your bus!

The Exponential distribution is often used to model *failure times* of components: for example $X \sim \text{Exponential}(\lambda)$ is the amount of time before a light bulb fails. In this case, memorylessness means that 'old is as good as new' — or, put another way, 'new is as bad as old'! A memoryless light bulb is quite likely to fail almost immediately.



For private reading: proof of memorylessness

nor examinable.

Let $X \sim \text{Exponential}(\lambda)$ be the *total* time waited for an event.

Let Y be the amount of *extra* time waited for the event, given that we have *already* waited time t (say).

We wish to prove that Y has the same distribution as X, i.e. that the time t already waited has been 'forgotten'. This means we need to prove that $Y \sim \text{Exponential}(\lambda)$.

<u>Proof</u>: We will work with $F_Y(y)$ and prove that it is equal to $1 - e^{-\lambda y}$. This proves that Y is Exponential(λ) like X.

First note that X = t + Y, because X is the *total* time waited, and Y is the time waited after time t. Also, we must condition on the event $\{X > t\}$, because we know that we have already waited time t. So $\mathbb{P}(Y \le y) = \mathbb{P}(X \le t + y | X > t)$.

$$F_Y(y) = \mathbb{P}(Y \le y) = \mathbb{P}(X \le t + y \mid X > t)$$
$$= \frac{\mathbb{P}(X \le t + y \quad \text{AND} \quad X > t)}{\mathbb{P}(X > t)}$$

(definition of conditional probability)

$$= \frac{\mathbb{P}(t < X \leq t + y)}{1 - \mathbb{P}(X \leq t)}$$

$$= \frac{F_X(t+y) - F_X(t)}{1 - F_X(t)}$$

$$= \frac{(1 - e^{-\lambda(t+y)}) - (1 - e^{-\lambda t})}{1 - (1 - e^{-\lambda t})}$$

$$= \frac{e^{-\lambda t} - e^{-\lambda(t+y)}}{e^{-\lambda t}}$$

$$= \frac{e^{-\lambda t}(1 - e^{-\lambda y})}{e^{-\lambda t}}$$

$$= 1 - e^{-\lambda y}. \quad \text{So } Y \sim \text{Exponential}(\lambda) \text{ as required.}$$

Thus the *conditional* probability of waiting time $y \ extra$, given that we have already waited time t, is the same as the probability of waiting time y in total. The time t already waited is forgotten.



4.4 Likelihood and estimation for continuous random variables

- For discrete random variables, we found the likelihood using the probability function, $f_{\mathbf{x}}(\mathbf{x}) = P(\mathbf{X}=\mathbf{x})$.
- For continuous random variables, we find the likelihood using the probability density function, $f_x(x) = \frac{dF_x}{dx}$
- Although the notation $f_X(x)$ means something different for continuous and discrete random variables, it is used in exactly the same way for likelihood and estimation.

Note: Both discrete and continuous r.v.<u>s</u> have the same definition for the cumulative distribution function: $f_{\chi}(\pi) = \mathbb{P}(\chi \leq \pi)$.

Example: Exponential likelihood

Suppose that:

- X~ Exponential (λ).
- · À is menown.
- · the observed value of X is x.

Then the likelihood function is:

$$\bot(\lambda;x)$$

We estimate
$$\lambda$$
 by setting $\frac{dL}{d\lambda} = 0$ to find the MLE, $\hat{\lambda}$.
or more independent observations $(Answo: \hat{\lambda} = \frac{L}{X})$.

Two or more independent observations

Suppose that X_1, \ldots, X_n are continuous random variables such that:

- X, ,..., X, are INDEPENDENT;
- all the Xi's have the same PDF, f (2);

then the likelihood is

$$L(\lambda; \pi_1, ..., \pi_n) = f_{\chi}(\pi_1) f_{\chi}(\pi_2) \cdots f_{\chi}(\pi_n),$$

for $0 < \lambda \leq \infty$.

le-xx for

0 <

Example: Suppose that X_1, X_2, \ldots, X_n are independent, and $X_i \sim \text{Exponential}(\lambda)$ for all *i*. Find the maximum likelihood estimate of λ .

the
$$\lambda$$
 is not guarding.
Likelihood graph shown
for $= 2$ and $n = 10$.
 $x_{1,...,x_{10}}$ generated
by R command
 $rexp(10, 2)$.
 $Solution:$
 $L(\lambda; \pi_{1},...,\pi_{n}) = \prod_{i=1}^{n} f_{X}(x_{i})$
 $= \prod_{i=1}^{n} (\lambda e^{-\lambda x_{i}})$
 $= \lambda e^{-\lambda(x_{1}+...+x_{n})}$
 $for 0 < \lambda < \infty$
Define $\bar{x} = \frac{1}{n} (\pi_{1}+...+\pi_{n})$, so $(x_{1}+...+x_{n}) = n\bar{x}$.
So $L(\lambda; \pi_{1},...,\pi_{n}) = \lambda e^{-n\bar{x}\lambda}$ for $0 < \lambda < \infty$.
For the MLE:
 $0 = dL = n\lambda^{n-1} e^{-n\bar{x}\lambda}$
 $1 = -\lambda\bar{z}$
 $= \lambda = 0$ or $\lambda = \infty$: not so because $0 < \lambda < \infty$
So the MLE is $\bar{\lambda} = \frac{1}{\bar{x}}$.

4.5 Hypothesis tests

P(X≤x) for p-value:

Hypothesis tests for continuous random variables are just like hypothesis tests for discrete random variables. The only difference is:

• Endpoints matter for discrete r.v.s. Endpoints DO NOT MATTER for continuous r.v.s.

Example: discrete. Suppose H_0 : $X \sim \text{Binomial}(n = 10, p = 0.5)$, and we have observed the value x = 7. Then the *upper-tail p*-value is

$$\mathbb{P}(X \ge 7) = 1 - \mathbb{P}(X \le 6) = 1 - f_{X}(6)$$

Discrete.

Example: continuous. Suppose H_0 : $X \sim \text{Exponential}(2)$, and we have observed the value x = 7. Then the upper-tail p-value is

 $P(X \ge 7) = I - P(X \le 7) = I - F_X(7).$ <u>CTS</u>.

Other than this trap, the procedure for hypothesis testing is the same:

- Use H_0 to specify the distribution of X completely, and offer a one-tailed or two-tailed alternative hypothesis H_1 .
- Make observation x.
- Find the one-tailed or two-tailed *p*-value as the probability of seeing an observation at least as weird as what we have seen, if H_0 is true.
- That is, find the probability under the distribution specified by H_0 of seeing an observation *further out in the tails* than the value x that we have seen.

Example with the Exponential distribution

A very very old person observes that the waiting time from Rangitoto to the next volcanic eruption in Auckland is 1500 years. Test the hypothesis that $\lambda = \frac{1}{1000}$ against the one-sided alternative that $\lambda < \frac{1}{1000}$.

Note: If $\lambda < \frac{1}{1000}$, we would expect to see LARGER values of X than would this is because X is the time between volcanoes, and λ is the rate at which volcanoes occur. A smaller value of λ means volcanoes occur LESS often, so the time waited, X, is LARGER.

OF AUCKLANE

<u>Hypotheses</u>: Let $X \sim E_{X\rho}$ (λ). H_{\circ} : $\lambda = \frac{1}{1000}$ H, : $\lambda < \frac{1}{1000}$ one-tailed test. **Observation:** x = 1500 years. Values weirder than x = 1500 years: all $\pi \ge 1500$ yrs, $P(X \ge 1500)$ when $X \sim E_{XP}(\frac{1}{1000})$. *p*-value: So p-value = $P(X \ge 1500)$ $= |-|P(X \leq 1500)$ $= |-F_{X}|(1500)$ $= 1 - \frac{1}{2} - e^{-\frac{1}{1000} * 1500}$ = 0.223. 1- pexp (1500, 1/1000). <u>*R* command:</u> Interpretation: There is no evidence Can't get widerce against the small A. Only the upportail gives wider a gainst against Ho. 1 2

The observation

$$\chi = 1500$$
 years
is compatible
with the hypothesis χ
ie. that orbitions occurs
on average.
 $\chi = 1000$
 $\chi = 10000$
 $\chi = 10000$
 $\chi = 10000$
 $\chi = 10000$
 $\chi = 1$

144

Read : use in Tut 10

4.6 Expectation and variance

Remember the expectation of a *discrete* random variable is *the long-term average:*

$$\mu_X = \mathbb{E}(X) = \sum_x x \mathbb{P}(X = x) = \sum_x x f_X(x).$$

(For each value x, we add in the value and multiply by the proportion of times we would expect to see that value: $\mathbb{P}(X = x)$.)

For a *continuous* random variable, replace the probability function with the probability density function, and replace \sum_x by $\int_{-\infty}^{\infty}$:

$$\mu_X = \mathbb{E}(X) = \int_{-\infty}^{\infty} x f_X(x) \, dx,$$

where $f_X(x) = F'_X(x)$ is the probability density function.

Note: There exists no concept of a 'probability function' $f_X(x) = \mathbb{P}(X = x)$ for continuous random variables. In fact, if X is continuous, then $\mathbb{P}(X = x) = 0$ for all x.

The idea behind expectation is the same for both discrete and continuous random variables. $\mathbb{E}(X)$ is:

- the long-term average of X;
- a 'sum' of values multiplied by how common they are: $\sum x f(x)$ or $\int x f(x) dx$.

Expectation is also the balance point of $f_X(x)$ for both continuous and discrete X. Imagine $f_X(x)$ cut out of cardboard and balanced on a pencil.

Z = Greek "S"

5-> [-> [

Discrete:

$$\mathbb{E}(X) = \sum_{x} x f_X(x)$$

$$\mathbb{E}(g(X)) = \sum_{x} g(x) f_X(x)$$

Transform the values, leave the probabilities alone;

$$f_X(x) = \mathbb{P}(X = x)$$

Continuous:

$$\mathbb{E}(X) = \int_{-\infty}^{\infty} x f_X(x) \, dx$$

$$\mathbb{E}(g(X)) = \int_{-\infty}^{\infty} g(x) f_X(x) \, dx$$

Transform the values, leave the probability density alone.

THE UNIVERSITY

$$f_X(x) = F'_X(x) \text{ (p.d.f.)}$$

Variance

If X is continuous, its variance is defined in exactly the same way as a discrete random variable:

$$Var(X) = \sigma_X^2 = \mathbb{E}\left((X - \mu_X)^2\right) = \mathbb{E}(X^2) - \mu_X^2 = \mathbb{E}(X^2) - (\mathbb{E}X)^2.$$

For a continuous random variable, we can either compute the variance using

$$\operatorname{Var}(X) = \mathbb{E}\left((X - \mu_X)^2\right) = \int_{-\infty}^{\infty} (x - \mu_X)^2 f_X(x) dx,$$

or

$$\operatorname{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}X)^2 = \int_{-\infty}^{\infty} x^2 f_X(x) dx - (\mathbb{E}X)^2.$$

The second expression is usually easier (although not always).

Properties of expectation and variance

All properties of expectation and variance are *exactly the same for continuous and discrete random variables.*

For any random variables, X, Y, and X_1, \ldots, X_n , continuous or discrete, and for constants a and b:

- $\mathbb{E}(aX+b) = a\mathbb{E}(X) + b.$
- $\mathbb{E}(ag(X) + b) = a\mathbb{E}(g(X)) + b.$
- $\mathbb{E}(X+Y) = \mathbb{E}(X) + \mathbb{E}(Y).$
- $\mathbb{E}(X_1 + \ldots + X_n) = \mathbb{E}(X_1) + \ldots + \mathbb{E}(X_n).$
- $\operatorname{Var}(aX + b) = a^2 \operatorname{Var}(X)$.
- $\operatorname{Var}(ag(X) + b) = a^2 \operatorname{Var}(g(X)).$

The following statements are generally true *only when X and Y are INDEPENDENT:*

- $\mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y)$ when X, Y independent.
- Var(X + Y) = Var(X) + Var(Y) when X, Y independent.

4.7 Exponential distribution mean and variance

Note: If X is the waiting time for a Poisson process with rate λ events per year (say), it makes sense that $\mathbb{E}(X) = \frac{1}{\lambda}$. For example, if $\lambda = 4$ events per hour, the average time waited between events is $\frac{1}{4}$ hour.

Know all Hris! **Proof:** $\mathbb{E}(X) = \int_{-\infty}^{\infty} x f_X(x) \, dx = \int_0^{\infty} x \lambda e^{-\lambda x} \, dx.$

No integ. by parts on Exam Tut O

Integration by parts: recall that
$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$
.
Let $u = x$, so $\frac{du}{dx} = 1$, and let $\frac{dv}{dx} = \lambda e^{-\lambda x}$, so $v = -e^{-\lambda x}$.
Then $\mathbb{E}(X) = \int_0^\infty x \lambda e^{-\lambda x} dx = \int_0^\infty u \frac{dv}{dx} dx$
 $= \left[uv \right]_0^\infty - \int_0^\infty v \frac{du}{dx} dx$
 $= \left[-xe^{-\lambda x} \right]_0^\infty - \int_0^\infty (-e^{-\lambda x}) dx$
 $= 0 + \left[\frac{-1}{\lambda} e^{-\lambda x} \right]_0^\infty$
 $= \frac{-1}{\lambda} \times 0 - \left(\frac{-1}{\lambda} \times e^0 \right)$
 $\therefore \mathbb{E}(X) = \frac{1}{\lambda}$.

<u>Variance</u>: $\operatorname{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}X)^2 = \mathbb{E}(X^2) - \frac{1}{\lambda^2}.$ Now $\mathbb{E}(X^2) = \int_{-\infty}^{\infty} x^2 f_X(x) \, dx = \int_0^{\infty} x^2 \lambda e^{-\lambda x} \, dx.$ Let $u = x^2$, so $\frac{du}{dx} = 2x$, and let $\frac{dv}{dx} = \lambda e^{-\lambda x}$, so $v = -e^{-\lambda x}.$ Then $\mathbb{E}(X^2) = \left[uv\right]_0^{\infty} - \int_0^{\infty} v \frac{du}{dx} \, dx = \left[-x^2 e^{-\lambda x}\right]_0^{\infty} + \int_0^{\infty} 2x e^{-\lambda x} \, dx$ $= 0 + \frac{2}{\lambda} \int_0^{\infty} \lambda x e^{-\lambda x} \, dx$ $= \frac{2}{\lambda} \times \mathbb{E}(X) = \frac{2}{\lambda^2}.$ So $\operatorname{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}X)^2 = \frac{2}{\lambda^2} - \left(\frac{1}{\lambda}\right)^2$ $\operatorname{Var}(X) = \frac{1}{\lambda^2}.$

Interlude: Guess the Mean, Median, and Variance

For any distribution:

- the *mean* is the *average* that would be obtained if a large number of observations were drawn from the distribution;
- the *median* is the *half-way point* of the distribution: every observation has a 50-50 chance of being above the median or below the median;
- the *variance* is the *average squared distance* of an observation from the mean.

Given the probability density function of a distribution, we should be able to guess roughly the distribution mean, median, and variance ... but it isn't easy! Have a go at the examples below. As a hint:

- the *mean* is the *balance-point* of the distribution. Imagine that the p.d.f. is made of cardboard and balanced on a rod. The mean is the point where the rod would have to be placed for the cardboard to balance.
- the *median* is the half-way point, so it divides the p.d.f. into two equal areas of 0.5 each.
- the *variance* is the average *squared* distance of observations from the mean; so to get a *rough* guess (not exact), it is easiest to guess an average distance from the mean and square it.

Answers:

Notes: The mean is larger than the median. This always happens when the distribution has a long right tail (positive skew) like this one.
The variance is *huge*... but when you look at the numbers along the horizontal axis, it is quite believable that the average squared distance of an observation from the mean is 118². Out of interest, the distribution shown is a Lognormal distribution.

Example 2: Try the same again with the example below. Answers are written below the graph.

4.8 The Uniform distribution

X has a Uniform distribution on the interval [a, b] if X is equally likely to fall anywhere in the interval [a, b]. We write X ~ Uniform [a, b], or X ~ Uniform (a, b) is equivalent. Or X ~ U[a, b] or X ~ U (a, b). Probability density function, $f_X(x)$

$$If X \sim U[a, b], then \qquad f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \le x \le b, \\ 0 & \text{otherwise.} \end{cases}$$

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \le x \le b, \\ 0 & \text{otherwise.} \end{cases}$$

$$f_X(x) = \int_{b-a}^{\infty} f_X(x)$$

$$f_X(x) = \int_{a}^{\infty} f_X(x) du$$

Mean and variance:

$$\mathbb{E}(X) = \int_{-\infty}^{\infty} xf(x) dx = \int_{a}^{b} x\left(\frac{1}{b-a}\right) dx = \frac{1}{b-a} \left[\frac{x^{2}}{2}\right]_{a}^{b}$$
$$= \left(\frac{1}{b-a}\right) \cdot \frac{1}{2}(b^{2}-a^{2})$$
$$= \left(\frac{1}{b-a}\right) \frac{1}{2}(b-a)(b+a)$$
$$= \frac{a+b}{2}.$$

Т

$$\begin{aligned} \operatorname{Var}(X) &= \mathbb{E}[(X - \mu_X)^2] = \int_a^b \frac{(x - \mu_X)^2}{b - a} \, dx \;\; = \;\; \frac{1}{b - a} \left[\frac{(x - \mu_X)^3}{3} \right]_a^b \\ &= \; \left(\frac{1}{b - a} \right) \left\{ \frac{(b - \mu_X)^3 - (a - \mu_X)^3}{3} \right\} \end{aligned}$$

But $\mu_X = \mathbb{E}X = \frac{a+b}{2}$, so $b - \mu_X = \frac{b-a}{2}$ and $a - \mu_X = \frac{a-b}{2}$. So,

$$\operatorname{Var}(X) = \left(\frac{1}{b-a}\right) \left\{ \frac{(b-a)^3 - (a-b)^3}{2^3 \times 3} \right\} = \frac{(b-a)^3 + (b-a)^3}{(b-a) \times 24}$$
$$= \frac{(b-a)^2}{12}. \square$$

Example: let $X \sim \text{Uniform}[0, 1]$. Then b - q = |- o = |

$$f_{X}(\pi) = \begin{cases} 1 & \text{for } 0 \le \pi \le 1 \\ 0 & \text{otherwise.} \end{cases}$$

and
$$EX = \frac{a+b}{2} = \frac{1}{2}$$

and $Var(X) = \frac{1}{12}(b-a)^2 = \frac{1}{12}$.

We know
$$P(a < X < b) = \int_{a}^{b} \frac{f_{x}(x)}{f_{x}(x)} dx$$

(defines the pdf)
What we need is $\# f_{y}(y)$ s.t.
 $P(c < Y < d) = \int_{c}^{d} \frac{f_{y}(y)}{f_{y}(y)} \frac{dy}{dy}$
Variable substitution in Calculus: $\left(\frac{f_{y}(y) - f_{x}(x(y))}{dy}\right)\frac{dz}{dy}$
e.g. integrate $\int_{0c}^{2^{2}} \sqrt{z} dx = \int_{0}^{2^{2}} \frac{y}{dy} \frac{dx}{dy}$
Let $y = \sqrt{x}$
Hen $x = y^{2}$
and $\frac{dx}{dy} = 2y$

Change of variable technique for monotone functions

Suppose that g(X) is a monotone function on the range of X. This means that g is an increasing function, or g is a decreasing function, or g is a decreasing function g is monotone, if is invotible, or (1-1) (one-to-one). That is, for every y, three is a UNIQUE π such that $g(\pi) = y$. This means that the inverse function, $g^{-1}(y)$, is well-defined as a function for a certain range of y.

When $g : \mathbb{R} \to \mathbb{R}$, as it is here, then g can only be (1–1) if it is monotone.

Change of Variable formula

Let $g : \mathbb{R} \to \mathbb{R}$ be a monotone function and let Y = g(X). Then the p.d.f. of Y = g(X) is

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$
 formal notation - ignore.

Easy way to remember

Write
$$y = y(x)$$
 $(=g(x))$
Then $x = x(y)$ $(=g^{-1}(y))$
Then $f_{y}(y) = f_{x}(x(y)) \left|\frac{dx}{dy}\right|$ LEARN

Working for change of variable questions

- 1) Show you have checked y = y(x) is monotone over the range of X.
- 2) Write $y = y(x) = \cdots$ for x in Krange of x)
- 3) Then $\chi = \chi(y) = \dots$ for y in (range of y)
- 4) Then $\left|\frac{dx}{dy}\right| = \langle differentiate #3 expr} \rangle$ $\longrightarrow expression involving y, not x.$

5) So
$$f_{y}(y) = f_{x}(x(y)) \left| \frac{dx}{dy} \right|$$
 by Change of Variable formula
= $\int_{-\infty}^{-\infty} \int_{0}^{0} f_{y}(y) dx + \int_{0}^{0} \int_{0}^{0} f_{y}(y) dx$

Quote range of values of y in your FINAL answer (1 mork). Refer back to the question to find $f_X(x)$: you often have to deduce this from information like $X \sim \text{Exponntial}(X)$ or $X \sim \text{Uniform}(0, 1)$. Or it may be given explicitly.

Note: There should be no x's left in the answer!

- 1. not stating g(x) is monotone over the required range of x;
- 2. not giving the range of y for which the result holds, as part of the FINAL answer. (eg. $f_Y(y) = \dots$ for $0 < y < \infty$).

Example 2: Let X be a continuous random variable with p.d.f.

$$f_{X}(x) = \begin{cases} \frac{1}{4}x^{3} & \text{for } 0 < x < 2, \\ 0 & \text{otherwise.} \end{cases}$$
Let $Y = 1/X$. Find the probability density function of Y , $f_{Y}(y)$.
1) $y = y(x) = \frac{1}{2}$ is monotone for $0 < x < 2$ => use C.O.V. formula.
2) $y = y(x) = \frac{1}{2}$ for $1 \le xy < \infty$.
3) $S_{0} = x = \frac{x}{(y)} = \frac{1}{y}$ for $\frac{1}{2} < y < \infty$.
4) $S_{0} = \left|\frac{4x}{4y}\right| = \left|-\frac{1}{y^{2}}\right| = \frac{1}{y^{2}}$ for $\frac{1}{2} < y < \infty$.
5) $S_{0} = f_{Y}(y) = f_{X}(x(y)) \left|\frac{4x}{4y}\right|$
 $= \frac{1}{4}\left(\frac{1}{y}\right)^{3} + \frac{1}{y^{2}}$ because
 $f_{X}(x) = \frac{1}{4}x^{3}$ in Q
 $\Rightarrow f_{Y}(y) = \frac{1}{4y^{5}}$ for $\frac{1}{2} < y < \infty$.
Find answer. (2)
(and O otherwise
is assemed if
you don't say
anything explicitly).

For mathematicians: proof of the change of variable formula

Separate into cases where g is increasing and where g is decreasing.

i) g increasing

g is increasing if $u < w \Leftrightarrow g(u) < g(w)$. \circledast Note that putting $u = g^{-1}(x)$, and $w = g^{-1}(y)$, we obtain

$$\begin{split} g^{-1}(x) < g^{-1}(y) & \Leftrightarrow \ g(g^{-1}(x)) < g(g^{-1}(y)) \\ & \Leftrightarrow \ x < y, \end{split}$$

so g^{-1} is also an increasing function.

Now

$$F_Y(y) = \mathbb{P}(Y \le y) = \mathbb{P}(g(X) \le y) = \mathbb{P}(X \le g^{-1}(y)) \quad \text{put} \begin{cases} u = X, \\ w = g^{-1}(y) \end{cases} \text{ in } \circledast \text{ to see this.} \\ = F_X(g^{-1}(y)). \end{cases}$$

So the p.d.f. of Y is

$$f_Y(y) = \frac{d}{dy} F_Y(y)$$

= $\frac{d}{dy} F_X(g^{-1}(y))$
= $F'_X(g^{-1}(y)) \frac{d}{dy}(g^{-1}(y))$ (Chain Rule)
= $f_X(g^{-1}(y)) \frac{d}{dy}(g^{-1}(y))$

Now g is increasing, so g^{-1} is also increasing (by overleaf), so $\frac{d}{dy}(g^{-1}(y)) > 0$, and thus $f_Y(y) = f_X(g^{-1}(y)) |\frac{d}{dy}(g^{-1}(y))|$ as required.

(Putting $u = g^{-1}(x)$ and $w = g^{-1}(y)$ gives $g^{-1}(x) > g^{-1}(y) \iff x < y$, so g^{-1} is also decreasing.)

$$F_Y(y) = \mathbb{P}(Y \le y) = \mathbb{P}(g(X) \le y)$$

= $\mathbb{P}(X \ge g^{-1}(y))$ (put $u = X, w = g^{-1}(y)$ in (\star))
= $1 - F_X(g^{-1}(y)).$

Thus the p.d.f. of Y is

$$f_Y(y) = \frac{d}{dy} \Big(1 - F_X(g^{-1}(y)) \Big) = -f_X \Big(g^{-1}(y) \Big) \frac{d}{dy} \Big(g^{-1}(y) \Big)$$

This time, g is decreasing, so g^{-1} is also decreasing, and thus

$$-\frac{d}{dy}\left(g^{-1}(y)\right) = \left|\frac{d}{dy}\left(g^{-1}(y)\right)\right|.$$

So once again,

$$f_Y(y) = f_X\left(g^{-1}(y)\right) \left| \frac{d}{dy}\left(g^{-1}(y)\right) \right|. \qquad \Box$$

4.10 Change of variable for non-monotone functions: non-examinable

Suppose that Y = g(X) and g is **not** monotone. We wish to find the p.d.f. of Y. We can sometimes do this by using the distribution function directly.

Example: Let X have **any** distribution, with distribution function $F_X(x)$. Let $Y = X^2$. Find the p.d.f. of Y.

Clearly,
$$Y \ge 0$$
, so $F_Y(y) = 0$ if $y < 0$.
For $y \ge 0$:
 $F_Y(y) = \mathbb{P}(Y \le y)$
 $= \mathbb{P}(X^2 \le y)$
 $= \mathbb{P}(-\sqrt{y} \le X \le \sqrt{y})$
 $= F_X(\sqrt{y}) - F_X(-\sqrt{y})$.

So

$$F_Y(y) = \begin{cases} 0 & \text{if } y < 0\\ F_X(\sqrt{y}) - F_X(-\sqrt{y}) & \text{if } y \ge 0 \end{cases}$$

So the p.d.f. of Y is

$$f_Y(y) = \frac{d}{dy} F_Y = \frac{d}{dy} (F_X(\sqrt{y})) - \frac{d}{dy} (F_X(-\sqrt{y}))$$
$$= \frac{1}{2} y^{-\frac{1}{2}} F'_X(\sqrt{y}) + \frac{1}{2} y^{-\frac{1}{2}} F'_X(-\sqrt{y})$$
$$= \frac{1}{2\sqrt{y}} \Big(f_X(\sqrt{y}) + f_X(-\sqrt{y}) \Big) \quad \text{for } y \ge 0.$$

$$\therefore \qquad f_Y(y) = \frac{1}{2\sqrt{y}} \left(f_X(\sqrt{y}) + f_X(-\sqrt{y}) \right) \text{ for } y \ge 0, \text{ whenever } Y = X^2.$$

Example: Let $X \sim \text{Normal}(0, 1)$. This is the familiar bell-shaped distribution (see later). The p.d.f. of X is:

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$

Find the p.d.f. of $Y = X^2$.

By the result above, $Y = X^2$ has p.d.f.

$$f_Y(y) = \frac{1}{2\sqrt{y}} \cdot \frac{1}{\sqrt{2\pi}} (e^{-y/2} + e^{-y/2})$$

= $\frac{1}{\sqrt{2\pi}} y^{-1/2} e^{-y/2}$ for $y \ge 0$.

This is in fact the Chi-squared distribution with $\nu = 1$ degrees of freedom.

The Chi-squared distribution is a special case of the Gamma distribution (see next section). This example has shown that if $X \sim \text{Normal}(0,1)$, then $Y = X^2 \sim \text{Chi-squared}(\text{df}=1)$.

4.11 The Gamma distribution

The Gamma (k, λ) distribution is a very flexible family of distributions.

It is defined as the sum of k independent Exponential r.v.s:

if $X_1, \ldots, X_k \sim \text{Exponential}(\lambda)$ and X_1, \ldots, X_k are independent, then $X_1 + X_2 + \ldots + X_k \sim \text{Gamma}(k, \lambda)$.

Special Case: When k = 1, $Gamma(1, \lambda) = Exponential(\lambda)$ (the sum of a single Exponential r.v.)

Probability density function, $f_X(x)$

For
$$X \sim \text{Gamma}(k, \lambda)$$
, $f_X(x) = \begin{cases} \frac{\lambda^k}{\Gamma(k)} x^{k-1} e^{-\lambda x} & \text{if } x \ge 0, \\ 0 & \text{otherwise.} \end{cases}$

Here, $\Gamma(k)$, called the Gamma function of k, is a constant that ensures $f_X(x)$ integrates to 1, i.e. $\int_0^\infty f_X(x) dx = 1$. It is defined as $\Gamma(k) = \int_0^\infty y^{k-1} e^{-y} dy$. When k is an integer, $\Gamma(k) = (k-1)!$

Mean and variance of the Gamma distribution:

For $X \sim \text{Gamma}(k, \lambda)$,

$$\mathbb{E}(X) = \frac{k}{\lambda}$$
 and $Var(X) = \frac{k}{\lambda^2}$

Relationship with the Chi-squared distribution

The Chi-squared distribution with ν degrees of freedom, χ^2_{ν} , is a special case of the Gamma distribution.

$$\chi^2_{\nu} = \operatorname{Gamma}(k = \frac{\nu}{2}, \lambda = \frac{1}{2}).$$

So if $Y \sim \chi^2_{\nu}$, then $\mathbb{E}(Y) = \frac{k}{\lambda} = \nu$, and $\operatorname{Var}(Y) = \frac{k}{\lambda^2} = 2\nu$.

Distribution function, $F_X(x)$

There is no closed form for the distribution function of the Gamma distribution. If $X \sim \text{Gamma}(k, \lambda)$, then $F_X(x)$ can can only be calculated by computer.

Proof that $\mathbb{E}(X) = \frac{k}{\lambda}$ and $\operatorname{Var}(X) = \frac{k}{\lambda^2}$ (non-examinable)

$$\mathbb{E}X = \int_0^\infty x f_X(x) \, dx = \int_0^\infty x \cdot \frac{\lambda^k x^{k-1}}{\Gamma(k)} e^{-\lambda x} \, dx$$

$$= \frac{\int_0^\infty (\lambda x)^k e^{-\lambda x} \, dx}{\Gamma(k)}$$

$$= \frac{\int_0^\infty y^k e^{-y}(\frac{1}{\lambda}) \, dy}{\Gamma(k)} \qquad (\text{letting } y = \lambda x, \ \frac{dx}{dy} = \frac{1}{\lambda})$$

$$= \frac{1}{\lambda} \cdot \frac{\Gamma(k+1)}{\Gamma(k)}$$

$$= \frac{1}{\lambda} \cdot \frac{k \Gamma(k)}{\Gamma(k)} \qquad (\text{property of the Gamma function}),$$

$$= \frac{k}{\lambda}.$$

$$\begin{aligned} \operatorname{Var}(X) &= \mathbb{E}(X^2) - (\mathbb{E}X)^2 = \int_0^\infty x^2 f_X(x) \, dx - \frac{k^2}{\lambda^2} \\ &= \int_0^\infty \frac{x^2 \lambda^k x^{k-1} e^{-\lambda x}}{\Gamma(k)} \, dx - \frac{k^2}{\lambda^2} \\ &= \frac{\int_0^\infty (\frac{1}{\lambda}) (\lambda x)^{k+1} e^{-\lambda x} \, dx}{\Gamma(k)} - \frac{k^2}{\lambda^2} \\ &= \frac{1}{\lambda^2} \cdot \frac{\int_0^\infty y^{k+1} e^{-y} \, dy}{\Gamma(k)} - \frac{k^2}{\lambda^2} \quad \left[\text{where } y = \lambda x, \frac{dx}{dy} = \frac{1}{\lambda} \right] \\ &= \frac{1}{\lambda^2} \cdot \frac{\Gamma(k+2)}{\Gamma(k)} - \frac{k^2}{\lambda^2} \\ &= \frac{1}{\lambda^2} \frac{(k+1)k \Gamma(k)}{\Gamma(k)} - \frac{k^2}{\lambda^2} \\ &= \frac{k}{\lambda^2}. \end{aligned}$$

Gamma distribution arising from the Poisson process

Recall that the waiting time between events in a Poisson process with rate λ has the *Exponential*(λ) *distribution*.

That is, if X_i =time waited between event i-1 and event i, then $X_i \sim Exp(\lambda)$.

The time waited from time 0 to the time of the kth event is

 $X_1 + X_2 + \ldots + X_k$, the sum of k independent Exponential(λ) r.v.s.

Thus the time waited until the kth event in a Poisson process with rate λ has the *Gamma* (k, λ) *distribution*.

Note: There are some similarities between the Exponential(λ) distribution and the (discrete) Geometric(p) distribution. Both distributions describe the 'waiting time' before an event. In the same way, the Gamma(k, λ) distribution is similar to the (discrete) Negative Binomial(k, p) distribution, as they both describe the 'waiting time' before the kth event.

4.12 The Beta Distribution: non-examinable

The Beta distribution has two parameters, α and β . We write $X \sim \text{Beta}(\alpha, \beta)$.

$$\begin{array}{c} \textbf{P.d.f.} \\ \hline \end{array} \qquad \qquad f(x) = \left\{ \begin{array}{cc} \frac{1}{B(\alpha,\beta)} x^{\alpha-1} (1-x)^{\beta-1} & \text{for } 0 < x < 1, \\ 0 & \text{otherwise.} \end{array} \right. \end{array}$$

The function $B(\alpha, \beta)$ is the *Beta function* and is defined by the integral

$$B(\alpha,\beta) = \int_0^1 x^{\alpha-1} (1-x)^{\beta-1} \, dx, \quad \text{for } \alpha > 0, \ \beta > 0.$$

It can be shown that $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)}.$