

Chapter 5: The Normal Distribution

and the Central Limit Theorem

The Normal distribution is the familiar bell-shaped distribution. It is probably the most important distribution in statistics, mainly because of its link with the Central Limit Theorem, which states that

Before studying the Central Limit Theorem, we look at the Normal distribution and some of its general properties.

5.1 The Normal Distribution

The Normal distribution has two parameters,

μ and σ^2 satisfy

We write


Probability density function, $f_X(x)$

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{\{-(x-\mu)^2/2\sigma^2\}} \quad \text{for } -\infty < x < \infty.$$


Distribution function, $F_X(x)$

There is no closed form for the distribution function of the Normal distribution. If $X \sim \text{Normal}(\mu, \sigma^2)$, then $F_X(x)$ can only be calculated
R command:

Probability density function, $f_X(x)$



Distribution function, $F_X(x)$



Mean and Variance

For $X \sim \text{Normal}(\mu, \sigma^2)$,

Linear transformations

If $X \sim \text{Normal}(\mu, \sigma^2)$, then for any constants a and b ,

$$aX + b \sim \text{Normal}(a\mu + b, a^2\sigma^2).$$

In particular,

$Z \sim \text{Normal}(0, 1)$ is called the

Proof that $aX + b \sim \text{Normal}(a\mu + b, a^2\sigma^2)$:

Let $X \sim \text{Normal}(\mu, \sigma^2)$, and let $Y = aX + b$. We wish to find the distribution of Y . Use *the change of variable technique*.

1) $y(x) = ax + b$ is monotone, so we can apply the Change of Variable technique.

2) Let $y = y(x) = ax + b$ for $-\infty < x < \infty$.

3) Then $x = x(y) = \frac{y-b}{a}$ for $-\infty < y < \infty$.

$$4) \left| \frac{dx}{dy} \right| = \left| \frac{1}{a} \right| = \frac{1}{|a|}.$$

$$5) \text{ So } f_Y(y) = f_X(x(y)) \left| \frac{dx}{dy} \right| = f_X\left(\frac{y-b}{a}\right) \frac{1}{|a|}. \quad (\star)$$

$$\text{But } X \sim \text{Normal}(\mu, \sigma^2), \text{ so } f_X(x) = \frac{1}{\sqrt{2\pi}\sigma^2} e^{-(x-\mu)^2/2\sigma^2}$$

$$\begin{aligned} \text{Thus } f_X\left(\frac{y-b}{a}\right) &= \frac{1}{\sqrt{2\pi}\sigma^2} e^{-\left(\frac{y-b}{a}-\mu\right)^2/2\sigma^2} \\ &= \frac{1}{\sqrt{2\pi}\sigma^2} e^{-(y-(a\mu+b))^2/2a^2\sigma^2}. \end{aligned}$$

Returning to (\star) ,

$$f_Y(y) = f_X\left(\frac{y-b}{a}\right) \cdot \frac{1}{|a|} = \frac{1}{\sqrt{2\pi}a^2\sigma^2} e^{-(y-(a\mu+b))^2/2a^2\sigma^2} \text{ for } -\infty < y < \infty.$$

But this is the p.d.f. of a $\text{Normal}(a\mu + b, a^2\sigma^2)$ random variable.

So, if $X \sim \text{Normal}(\mu, \sigma^2)$, then $aX + b \sim \text{Normal}(a\mu + b, a^2\sigma^2)$.

Sums of Normal random variables

If X and Y are and $X \sim \text{Normal}(\mu_1, \sigma_1^2)$, $Y \sim \text{Normal}(\mu_2, \sigma_2^2)$,
then

More generally, if X_1, X_2, \dots, X_n are , and $X_i \sim \text{Normal}(\mu_i, \sigma_i^2)$
for $i = 1, \dots, n$, then

$$a_1 X_1 + a_2 X_2 + \dots + a_n X_n \sim \text{Normal}\left((a_1 \mu_1 + \dots + a_n \mu_n), (a_1^2 \sigma_1^2 + \dots + a_n^2 \sigma_n^2)\right).$$

For mathematicians: properties of the Normal distribution

1. Proof that $\int_{-\infty}^{\infty} f_X(x) dx = 1$.

The full proof that $\int_{-\infty}^{\infty} f_X(x) dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}\sigma^2} e^{\{-(x-\mu)^2/(2\sigma^2)\}} dx = 1$

relies on the following result:

$$\text{FACT: } \int_{-\infty}^{\infty} e^{-y^2} dy = \sqrt{\pi}.$$

This result is non-trivial to prove. See Calculus courses for details.

Using this result, the proof that $\int_{-\infty}^{\infty} f_X(x) dx = 1$ follows by using the change of variable $y = \frac{(x-\mu)}{\sqrt{2}\sigma}$ in the integral.

2. Proof that $\mathbb{E}(X) = \mu$.

$$\mathbb{E}(X) = \int_{-\infty}^{\infty} x f_X(x) dx = \int_{-\infty}^{\infty} x \frac{1}{\sqrt{2\pi}\sigma^2} e^{-(x-\mu)^2/2\sigma^2} dx$$

Change variable of integration: let $z = \frac{x-\mu}{\sigma}$: then $x = \sigma z + \mu$ and $\frac{dx}{dz} = \sigma$.

$$\text{Then } \mathbb{E}(X) = \int_{-\infty}^{\infty} (\sigma z + \mu) \cdot \frac{1}{\sqrt{2\pi}\sigma^2} \cdot e^{-z^2/2} \cdot \sigma dz$$

$$= \underbrace{\int_{-\infty}^{\infty} \frac{\sigma z}{\sqrt{2\pi}} \cdot e^{-z^2/2} dz}_{\text{this is an odd function of } z \text{ (i.e. } g(-z) = -g(z)\text{), so it integrates to 0 over range } -\infty \text{ to } \infty.} + \mu \underbrace{\int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz}_{\text{p.d.f. of } N(0,1) \text{ integrates to 1.}}$$

$$\begin{aligned} \text{Thus } \mathbb{E}(X) &= 0 + \mu \times 1 \\ &= \mu. \end{aligned}$$


3. Proof that $\text{Var}(X) = \sigma^2$.

$$\begin{aligned} \text{Var}(X) &= E \{ (X - \mu)^2 \} \\ &= \int_{-\infty}^{\infty} (x - \mu)^2 \frac{1}{\sqrt{2\pi}\sigma^2} e^{-(x-\mu)^2/(2\sigma^2)} dx \\ &= \sigma^2 \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} z^2 e^{-z^2/2} dz \quad \left(\text{putting } z = \frac{x - \mu}{\sigma} \right) \\ &= \sigma^2 \left\{ \frac{1}{\sqrt{2\pi}} \left[-ze^{-z^2/2} \right]_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz \right\} \quad (\text{integration by parts}) \\ &= \sigma^2 \{0 + 1\} \\ &= \sigma^2. \end{aligned}$$

□

5.2 The Central Limit Theorem (CLT)

also known as... the Piece of Cake Theorem



The Central Limit Theorem (CLT) is one of the most fundamental results in statistics. In its simplest form, it states that if a large number of independent random variables are drawn from **any** distribution, then the distribution of their sum (or alternatively their sample average) always converges to the Normal distribution.

Distribution of the sample mean, \bar{X} , using the CLT

Let X_1, \dots, X_n be independent, identically distributed with mean μ and variance σ^2 for all i .

The sample mean, \bar{X} , is defined as:

So $\bar{X} = \frac{S_n}{n}$, where

Because \bar{X} is a scalar multiple of a Normal r.v. as n grows large, \bar{X} *itself is approximately Normal for large n* :

The following three statements of the Central Limit Theorem are equivalent:

$$\begin{aligned}\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n} &\sim \text{approx Normal} \left(\mu, \frac{\sigma^2}{n} \right) \text{ as } n \rightarrow \infty. \\ S_n = X_1 + X_2 + \dots + X_n &\sim \text{approx Normal} (n\mu, n\sigma^2) \text{ as } n \rightarrow \infty. \\ \frac{S_n - n\mu}{\sqrt{n\sigma^2}} = \frac{\bar{X} - \mu}{\sqrt{\sigma^2/n}} &\sim \text{approx Normal} (0, 1) \text{ as } n \rightarrow \infty.\end{aligned}$$

The essential point to remember about the Central Limit Theorem is that large sums or sample means of independent random variables converge to a Normal distribution,

More general version of the CLT

A more general form of CLT states that, if X_1, \dots, X_n are independent, and $\mathbb{E}(X_i) = \mu_i$, $\text{Var}(X_i) = \sigma_i^2$ (not necessarily all equal), then

$$Z_n = \frac{\sum_{i=1}^n (X_i - \mu_i)}{\sqrt{\sum_{i=1}^n \sigma_i^2}} \rightarrow \text{Normal}(0, 1) \text{ as } n \rightarrow \infty.$$


Other versions of the CLT relax the condition that X_1, \dots, X_n are independent.

The Central Limit Theorem in action : simulation studies

The following simulation study illustrates the Central Limit Theorem, making use of several of the techniques learnt in STATS 210. We will look particularly at *how fast the distribution of S_n converges to the Normal distribution*.

Example 1: Triangular distribution: $f_X(x) = 2x$ for $0 < x < 1$.

Find $\mathbb{E}(X)$ and $\text{Var}(X)$:




Let $S_n = X_1 + \dots + X_n$ where X_1, \dots, X_n are


Then

So

The graph shows histograms of 10 000 values of $S_n = X_1 + \dots + X_n$ for $n = 1, 2, 3$, and 10. The Normal p.d.f. $\text{Normal}(n\mu, n\sigma^2) = \text{Normal}(\frac{2n}{3}, \frac{n}{18})$ is superimposed across the top. Even for n as low as 10, the Normal curve is a very good approximation.



Example 2: U-shaped distribution: $f_X(x) = \frac{3}{2}x^2$ for $-1 < x < 1$.




We find that

Let $S_n = X_1 + \dots + X_n$ where X_1, \dots, X_n are

Then

So

Even with this highly non-Normal distribution for X , the Normal curve provides a good approximation to $S_n = X_1 + \dots + X_n$ for n as small as 10.



Normal approximation to the Binomial distribution, using the CLT

Let $Y \sim \text{Binomial}(n, p)$.

We can think of Y as the *sum of n Bernoulli random variables*:

$Y = X_1 + X_2 + \dots + X_n$, where

Thus by the CLT,

Thus,


$$\text{Bin}(n, p) \rightarrow \text{Normal}\left(\underbrace{np}_{\text{mean of Bin}(n,p)}, \underbrace{np(1-p)}_{\text{var of Bin}(n,p)}\right) \text{ as } n \rightarrow \infty \text{ with } p \text{ fixed.}$$

The Binomial distribution is therefore well approximated by the Normal distribution when n is large, for any fixed value of p .


The Normal distribution is also a good approximation to the Poisson(λ) distribution when λ is large:

$$\text{Poisson}(\lambda) \rightarrow \text{Normal}(\lambda, \lambda) \text{ when } \lambda \text{ is large.}$$

Binomial($n = 100, p = 0.5$)



Poisson($\lambda = 100$)



Why the Piece of Cake Theorem? ...



- The Central Limit Theorem makes whole realms of statistics into a
 - After seeing a theorem this good, you deserve
-

5.3 Confidence intervals

Example: Remember the margin of error for an opinion poll?

An opinion pollster wishes to estimate the level of support for Labour in an upcoming election. She interviews n people about their voting preferences. Let p be the true, unknown level of support for the Labour party in New Zealand. Let X be the number of of the n people interviewed by the opinion pollster who plan to vote Labour. Then

At the end of Chapter 2, we said that the maximum likelihood estimator for p is

In a large sample (large n), we now know that


So

So

Now if $Z \sim \text{Normal}(0, 1)$, we find (using a computer) that the 95% central probability region of Z is from -1.96 to $+1.96$:

$$\mathbb{P}(-1.96 < Z < 1.96) = 0.95.$$

Check in *R*: `pnorm(1.96, mean=0, sd=1) - pnorm(-1.96, mean=0, sd=1)`



Putting $Z = \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}}$, we obtain

$$\mathbb{P} \left(-1.96 < \frac{\hat{p} - p}{\sqrt{\frac{pq}{n}}} < 1.96 \right) \simeq 0.95.$$

Rearranging

$$\mathbb{P} \left(\hat{p} - 1.96\sqrt{\frac{pq}{n}} < p < \hat{p} + 1.96\sqrt{\frac{pq}{n}} \right) \simeq 0.95.$$

This enables us to form an estimated 95% confidence interval for the unknown parameter p :

The 95% confidence interval *has RANDOM end-points, which depend on \hat{p} . About 95% of the time, these random end-points will enclose the true unknown value, p .*

Confidence intervals are extremely important for helping us to assess

A narrow confidence interval suggests

A wide confidence interval suggests

When you see newspapers quoting the *margin of error* on an opinion poll:

- Remember:
- Think:
- Have:

Confidence intervals for the Poisson λ parameter

We saw in section 3.6 that if X_1, \dots, X_n are independent, identically distributed with $X_i \sim \text{Poisson}(\lambda)$, then the maximum likelihood estimator of λ is

$$\hat{\lambda} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i.$$

Now

Thus, when n is large,

$$\hat{\lambda} = \bar{X} \sim \text{approx Normal}(\mu, \frac{\sigma^2}{n})$$

by the Central Limit Theorem. In other words,

We use the same transformation as before to find approximate 95% confidence intervals for λ as n grows large:

Let $Z = \frac{\hat{\lambda} - \lambda}{\sqrt{\frac{\lambda}{n}}}$. We have

Thus:

$$\mathbb{P} \left(-1.96 < \frac{\hat{\lambda} - \lambda}{\sqrt{\frac{\lambda}{n}}} < 1.96 \right) \simeq 0.95.$$

Rearranging

$$\mathbb{P} \left(\hat{\lambda} - 1.96\sqrt{\frac{\lambda}{n}} < \lambda < \hat{\lambda} + 1.96\sqrt{\frac{\lambda}{n}} \right) \simeq 0.95.$$

So our **estimated** 95% confidence interval for the unknown parameter λ is:

Why is this so good?

It's clear that it's important to measure precision, or reliability, of an estimate, otherwise the estimate is almost worthless. However, we have already seen various measures of precision: variance, standard error, coefficient of variation, and now confidence intervals. Why do we need so many?

- The true variance of an estimator, e.g. $\text{Var}(\hat{\lambda})$, is the most convenient quantity to work with mathematically. However, it is on a non-intuitive scale (squared deviation from the mean), and it usually depends upon the unknown parameter, e.g. λ .
- The **standard error** is $\text{se}(\hat{\lambda}) = \sqrt{\widehat{\text{Var}}(\hat{\lambda})}$. It is an **estimate** of the square root of the true variance, $\text{Var}(\hat{\lambda})$. Because of the square root, the standard error is a direct measure of deviation from the mean, rather than squared deviation from the mean. This means it is measured in more intuitive units. However, it is still unclear how we should comprehend the information that the standard error gives us.
- The beauty of the Central Limit Theorem is that it gives us an incredibly easy way of understanding what the standard error is telling us, using **Normal-based asymptotic confidence intervals** as computed in the previous two examples.

Although it is beyond the scope of this course to see why, the Central Limit Theorem guarantees that almost **any** maximum likelihood estimator will be Normally distributed as long as the sample size n is large enough, subject only to fairly mild conditions.

Thus,

or equivalently,

$$\hat{\lambda} - 1.96 \text{se}(\hat{\lambda}) \quad \text{to} \quad \hat{\lambda} + 1.96 \text{se}(\hat{\lambda}) .$$

The confidence interval has an easily-understood interpretation:

So the Central Limit Theorem has given us an incredibly simple and powerful way of converting from a hard-to-understand measure of precision, $\text{se}(\hat{\lambda})$, to a measure that is easily understood and relevant to the problem at hand. Brilliant!