

# THE UNIVERSITY OF AUCKLAND

---

SECOND SEMESTER, 2012

Campus: City

---

## STATISTICS


Special Topic in Applied Probability

(Time allowed: THREE hours)

**NOTE:** Attempt **ALL** questions. Marks for each question are shown in brackets.  
There are 100 marks in total.  
An **Attachment** containing useful information is found on page 8.

CONTINUED

1. A knight on an infinitely large chessboard always has eight different moves available to it.



The knight walks randomly on the board: at each step, a move is chosen at random, independently of anything else, with the eight available moves each being chosen with probability  $1/8$ . Call the starting square  $(0, 0)$ , and let  $(X_t, Y_t)$  be the square the knight has reached after  $t$  moves. Let  $R_t = \sqrt{X_t^2 + Y_t^2}$ , the knight's distance from its starting square after  $t$  moves.

- (a) Find a constant  $\beta$  such that

$$M_t = R_t^2 - \beta t$$

is a martingale with respect to  $(X_t, Y_t)$ .

(4)

- (b) For each time  $t$ , find the maximum and minimum possible values, and also the expected value, of  $R_t^2$ .

(3)

- (c) Let  $r > 0$  be a constant, and let  $T$  be the first time  $t$  with  $R_t \geq r$ . Explain why  $T$  is a stopping time with respect to  $(X_t, Y_t)$ .

(3)

- (d) Explain why  $E[T] < \infty$ .

(4)

- (e) Find  $E[M_T]$ , and from this show that

$$E[T] \geq \frac{r^2}{5}.$$

Carefully justify any assumptions you make.

(4)

[18 marks]

2. A test for a particular disease operates as follows. The patient provides a blood sample, which goes through a first test. If the test shows that the patient is infected with the disease, the blood sample is sent for a second test to diagnose cause and severity.


The first test takes  $3 + X$  days to complete, where  $X \sim \text{Poisson}(1)$ . The second test takes a further  $Y$  days to complete, where  $Y \sim \text{Poisson}(2)$ . Assume that  $X$  and  $Y$  are independent.

Define the indicator random variable  $D$ , such that

$$D = \begin{cases} 1 & \text{if patient has the disease,} \\ 0 & \text{if patient is healthy.} \end{cases}$$

Suppose that  $\mathbb{P}(D = 1) = 0.2$  in the population being tested.

A patient's progress through the testing process is shown in the diagram below.




Define the random variable  $T$  to be the total time taken before the sample testing is complete. Thus  $T$  is the total time taken for the first test and (if needed) the second test.

- (a) Show that the total expected testing time is  $\mathbb{E}(T) = 4.4$  days. You may use any results from the Attachment that you need. (2)
- (b) The probability generating functions (PGFs) of  $X$  and  $Y$  are  $G_X(s) = \mathbb{E}(s^X) = e^{s-1}$  for  $X \sim \text{Poisson}(1)$ , and  $G_Y(s) = \mathbb{E}(s^Y) = e^{2(s-1)}$  for  $Y \sim \text{Poisson}(2)$ . Find  $G_{X+Y}(s)$ , the PGF of  $X + Y$ , and hence name the distribution of  $X + Y$ , giving the values of any parameters. (3)
- (c) Find  $\mathbb{P}(T \geq 5 | D = 0)$ , the probability that a healthy sample will take at least 5 days to process; and  $\mathbb{P}(T \geq 5 | D = 1)$ , the probability that a diseased sample will take at least 5 days to process. (3)
- (d) An anxious patient has already waited 5 days for her test result. She wants to know the probability that she has the disease, given that her sample has taken at least 5 days to process. Find  $\mathbb{P}(D = 1 | T \geq 5)$ . (2)
- (e) Using the law of total variance, or otherwise, find  $\text{Var}(T)$ . (5)

[15 marks]

3. Let  $\{X_0, X_1, X_2, \dots\}$  be a random walk on the integers, with transition diagram below.



Let  $U$  be the number of steps taken to reach state 1, starting at state 0. Let  $H_U(s) = \mathbb{E}(s^U)$  be the probability generating function of  $U$ .

(a) Show that  $H_U(s)$  must be either the (+) root or the (-) root of the following expression:

$$H_U(s) = \frac{4 \pm \sqrt{16 - 12s^2}}{2s}. \tag{4}$$

(b) By considering  $\lim_{s \rightarrow 0} H_U(s)$ , prove that  $H_U(s)$  can **not** be the (+) root in the expression above. (2)


(c) Using the expression  $H_U(s) = \frac{4 - \sqrt{16 - 12s^2}}{2s}$ , find whether  $U$  is a defective random variable. Hence state the probability that the process *never* reaches state 1, starting from state 0. (3)

Let  $V$  be the number of steps taken to reach state  $-1$ , starting at state 0. Let  $H_V(s) = \mathbb{E}(s^V)$  be the probability generating function of  $V$ . You may assume that

$$H_V(s) = \frac{4 - \sqrt{16 - 12s^2}}{6s}.$$

(d) Let  $T$  be the number of steps taken to **first return** to state 0, starting at state 0. For example, if  $X_0 = 0, X_1 = 1,$  and  $X_2 = 0,$  then  $T = 2$  steps are taken to return from 0 to 0 again. Let  $G(s) = \mathbb{E}(s^T)$  be the probability generating function of  $T$ . Using  $G(s)$ , find the probability that the process *never* returns to state 0, starting from state 0. (3)

(e) Now suppose that the random walk has a boundary at state 100, as shown in the transition diagram below. Are your calculations for the probabilities in parts (c) and (d) still valid? If not, briefly explain how those probabilities could be correctly calculated. (4)



[16 marks]

4. A man owns  $N$  umbrellas. Every day, he walks between his home and his workplace and back again. On each of his trips, from home to work or from work to home, he carries an umbrella only if he has an umbrella available and it is raining. It rains on each trip with probability  $r$ , independently of all other trips, where  $0 < r < 1$ .

Let  $\{X_1, X_2, X_3, \dots\}$  be a Markov chain on the state space  $S = \{0, 1, 2, \dots, N\}$ , such that  $X_t$  is the **number of umbrellas available** on trip  $t$ . For example, if trip  $t$  is a trip from home to work, then  $X_t = n$  if there are  $n$  umbrellas at home, with the other  $N - n$  umbrellas being at work.

- (a) Draw the transition diagram when the man owns  $N = 1$  umbrella. Write down the equilibrium equations, and solve them to find the equilibrium distribution  $\boldsymbol{\pi}^T = (\pi_0, \pi_1)$  when  $N = 1$ . (4)
- (b) Draw the transition diagram when the man owns  $N = 2$  umbrellas. By direct substitution into the appropriate equilibrium equations, verify that the Markov chain has the following equilibrium distribution when  $N = 2$ :

$$\boldsymbol{\pi}^T = (\pi_0, \pi_1, \pi_2) = \left( \frac{1-r}{3-r}, \frac{1}{3-r}, \frac{1}{3-r} \right). \quad (4)$$

- (c) Suppose the man has  $N$  umbrellas. We wish to prove that the Markov chain has the following equilibrium distribution:

$$\boldsymbol{\pi}_*^T = (\pi_0, \pi_1, \pi_2, \dots, \pi_N) = \frac{1}{N+1-r} (1-r, 1, 1, \dots, 1).$$

Find  $\mathbb{E}(s^{X_{t+1}} | X_t)$ , and hence show that  $X_{t+1} \sim X_t$  if  $X_t \sim \boldsymbol{\pi}_*$ . Deduce that  $\boldsymbol{\pi}_*$  is the equilibrium distribution for this Markov chain, and (with justification) state the long-term probability that the man gets wet on a trip. You may assume that the chain is aperiodic for any value of  $N$ . (6)

[14 marks]

5. Let  $\{X_1, X_2, X_3, \dots\}$  be a Markov chain on the state space  $S = \{1, 2, \dots, N\}$ , with transition matrix  $P = (p_{ij})$ , such that:

- for  $i = 2, 3, \dots, N$ ,

$$p_{ij} = \begin{cases} \frac{1}{i} & \text{for } j = 1, \dots, i, \\ 0 & \text{for } j = i + 1, \dots, N; \end{cases}$$

- for  $i = 1$ ,

$$p_{1j} = \begin{cases} 0 & \text{for } j = 1, \dots, N - 1, \\ 1 & \text{for } j = N. \end{cases}$$

(a) Draw the transition diagram for  $N = 4$ . (2)

(b) Does the Markov chain  $\{X_t\}_{t \geq 1}$  converge to an equilibrium distribution as  $t \rightarrow \infty$ , for any  $N \geq 2$ ? Explain why or why not. (2)

(c) For  $x = 1, 2, \dots, N$ , define  $m_x$  to be the expected reaching time for state 1, starting from state  $x$ : that is,

$$m_x = \mathbb{E}(\text{number of steps taken to hit state 1} \mid \text{start at state } x).$$

We define  $m_1 = 0$ . Using first-step analysis, show that

$$m_{x+1} = m_x + \frac{1}{x} \quad \text{for } x = 2, \dots, N - 1,$$

and find  $m_2$ . (4)

(d) A theorem that we have not studied states that, for an irreducible Markov chain on a finite state space, the equilibrium distribution  $\pi$  satisfies

$$\pi_k = \frac{1}{R_{kk}}$$

where  $R_{kk}$  is the **expected return time** for state  $k$ . That is,  $R_{kk}$  is the expected number of steps needed to return to state  $k$  for the first time, starting at state  $k$ .

(Note that  $R_{kk} \geq 1$ , because at least one step is needed to *return* to state  $k$ , starting from state  $k$ . The expected return times differ from the expected reaching times, which are defined as 0 when starting at the target state.)

Show that the long-run proportion of time that the Markov chain spends in state 1 is

$$\pi_1 = \left( 2 + \sum_{r=1}^{N-1} \frac{1}{r} \right)^{-1}.$$

You must fully justify all parts of your answer. (4)

[12 marks]

6. Let  $\{Z_0, Z_1, Z_2, \dots\}$  be a branching process, where  $Z_n$  denotes the population size at time  $n$ , and  $Z_0 = 1$ . Let  $Y$  be the family size distribution. Suppose that  $Y \sim \text{Geometric}(p = 0.5)$ , so that

$$\mathbb{P}(Y = y) = \left(\frac{1}{2}\right)^{y+1} \quad \text{for } y = 0, 1, 2, \dots$$

- (a) Let  $G(s) = \mathbb{E}(s^Y)$  be the probability generating function of  $Y$ . Show that

$$G(s) = \frac{1}{2-s},$$

and state the range of values of  $s$  for which this expression is valid. (3)

- (b) Find the probability of eventual extinction,  $\gamma$ . (3)

- (c) Let  $G_n(s) = \mathbb{E}(s^{Z_n})$  be the probability generating function of the population size at time  $n$ . State (without calculation) the correct expression for  $G_{n+1}(s)$  in terms of  $G_n(s)$ . That is, rewrite the expression below, with  $\star$  replaced by the correct quantity:

$$G_{n+1}(s) = G_n(\star). \tag{1}$$

- (d) For  $Y \sim \text{Geometric}(p = 0.5)$  as above, prove by mathematical induction that

$$G_n(s) = \frac{n - (n-1)s}{(n+1) - ns} \quad \text{for } n = 1, 2, 3, \dots \tag{6}$$

- (e) Using the expression shown in (d), state expressions for the following probabilities:

- (i)  $\gamma_n$ , that the process is extinct by generation  $n$ : that is,  $\gamma_n = \mathbb{P}(Z_n = 0)$ ;
- (ii)  $\rho_n$ , that there are surviving individuals in generation  $n$ : that is,  $\rho_n = \mathbb{P}(Z_n > 0)$ . (2)

- (f) An idea of interest in branching processes is the **Most Recent Common Ancestor**. If there are  $z$  individuals alive in generation  $n$ , we can trace back the parents, grandparents, and so on, for each of the  $z$  individuals until we get to a single individual in an earlier generation who is a common ancestor for all  $z$  individuals. The **most recent** common ancestor is the most recent of all common ancestors: that is, the common ancestor that lived closest in time to generation  $n$ .

Let  $T_n$  be the generation number of the most recent common ancestor for individuals surviving at generation  $n$ . For example, if all individuals at time  $n$  had the same parent at time  $n - 1$ , then  $T_n = n - 1$ . Alternatively, if there is no common ancestor until the single individual at generation 0, then  $T_n = 0$ . (Note that the individual who started the branching process at time 0 is always a common ancestor to all individuals alive at generation  $n$ .)

For the branching process above, with  $Y \sim \text{Geometric}(p = 0.5)$ , show that

$$\mathbb{P}(Z_n > 0 \text{ and } T_n = 0) = \sum_{y=2}^{\infty} \left(\frac{1}{2}\right)^{y+1} \left\{ 1 - \left(\frac{n-1}{n}\right)^y - y \left(\frac{n-1}{n}\right)^{y-1} \left(\frac{1}{n}\right) \right\}.$$

Hence find an expression for  $\alpha_{n,0}$ : the conditional probability, given that there are individuals still alive in generation  $n$ , that their most recent common ancestor lived in generation 0.

Lastly, find a similar expression for  $\alpha_{n,1}$  in terms of  $\alpha_{n-1,0}$ , where  $\alpha_{n,1} = \mathbb{P}(T_n = 1 \mid Z_n > 0)$ . (10)

[25 marks]

## ATTACHMENT

## 1. Discrete Probability Distributions

Distribution	$\mathbb{P}(X = x)$	$\mathbb{E}(X)$	$\text{Var}(X)$	PGF, $\mathbb{E}(s^X)$
Geometric( $p$ )	$pq^x$ (where $q = 1 - p$ ), for $x = 0, 1, 2, \dots$	$\frac{q}{p}$	$\frac{q}{p^2}$	$\frac{p}{1 - qs}$
	Number of failures before the first success in a sequence of independent trials, each with $\mathbb{P}(\text{success}) = p$ .			
Binomial( $n, p$ )	$\binom{n}{x} p^x q^{n-x}$ (where $q = 1 - p$ ), for $x = 0, 1, 2, \dots, n$ .	$np$	$npq$	$(ps + q)^n$
	Number of successes in $n$ independent trials, each with $\mathbb{P}(\text{success}) = p$ .			
Poisson( $\lambda$ )	$\frac{\lambda^x}{x!} e^{-\lambda}$ for $x = 0, 1, 2, \dots$	$\lambda$	$\lambda$	$e^{\lambda(s-1)}$

2. Uniform Distribution:  $X \sim \text{Uniform}(a, b)$ .

Probability density function,  $f_X(x) = \frac{1}{b-a}$  for  $a < x < b$ . Mean,  $\mathbb{E}(X) = \frac{a+b}{2}$ .

## 3. Properties of Probability Generating Functions

**Definition:**  $G_X(s) = \mathbb{E}(s^X)$

**Moments:**  $\mathbb{E}(X) = G'_X(1)$        $\mathbb{E}\left\{X(X-1)\dots(X-k+1)\right\} = G_X^{(k)}(1)$

**Probabilities:**  $\mathbb{P}(X = n) = \frac{1}{n!} G_X^{(n)}(0)$

4. Geometric Series:  $1 + r + r^2 + r^3 + \dots = \sum_{x=0}^{\infty} r^x = \frac{1}{1-r}$  for  $|r| < 1$ .

Finite sum:  $\sum_{x=0}^n r^x = \frac{1-r^{n+1}}{1-r}$  for  $r \neq 1$ .

5. Binomial Theorem: For any  $p, q \in \mathbb{R}$ , and integer  $n > 0$ ,  $(p+q)^n = \sum_{x=0}^n \binom{n}{x} p^x q^{n-x}$ .6. Exponential Power Series: For any  $\lambda \in \mathbb{R}$ ,  $\sum_{x=0}^{\infty} \frac{\lambda^x}{x!} = e^\lambda$ .