

Chapter 7: Extinction in Branching Processes



Revision: a branching process consists of reproducing individuals.

- All individuals are independent.
- Start with a single individual at time 0: $Z_0 = 1$.
- Each individual lives a single unit of time, then has Y offspring and dies.
- Let Z_n be the size of generation n : the number of individuals born at time n .
- The branching process is $\{Z_0 = 1, Z_1, Z_2, \dots\}$.

Branching Process Recursion Formula

This is the fundamental formula for branching processes. Let $G_n(s) = \mathbb{E}(s^{Z_n})$ be the PGF of Z_n , the population size at time n . Let $G(s) = G_1(s)$, the PGF of the family size distribution Y , or equivalently, of Z_1 . Then:

$$G_n(s) = G_{n-1}(G(s)) = \underbrace{G(G(G(\dots G(s)\dots)))}_{n \text{ times}} = G(G_{n-1}(s)).$$

$$G_n(s) = \mathbb{E}(s^{Z_n})$$

$$G(s) = \mathbb{E}(s^Y) = \mathbb{E}(s^{Z_1})$$

✓ $P(\text{eventually goes extinct?})$

It is possible to find several results about the probability of eventual extinction.

If $Z_n = 0$, then the population is extinct for ever:
 $Z_t = 0$ for all $t \geq n$.



$$E_n = \{Z_n = 0\} = \{\text{extinct BY generation } n\}.$$

$$E_0 \subseteq E_1 \subseteq E_2 \subseteq E_3 \subseteq E_4 \subseteq \dots$$

This is because event E_i forces E_j to be true for all $j \geq i$, so E_i is a 'part' or subset of E_j for $j \geq i$. *increasing series of events.*

At the start of the branching process, we are interested in the probability of ultimate extinction: what is the probability of extinction at any point n in the future?

We can express this probability in different ways:

$$\mathbb{P}(\text{ultimate extinction}) = \mathbb{P}\left(\begin{matrix} \text{extinct by gen } 0 & \text{or} \\ \text{"} & \text{"} & \text{"} & \text{1} & \text{or} \\ \text{"} & \text{"} & \text{"} & \text{2} & \text{or} \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{matrix}\right) = \mathbb{P}\left(\bigcup_{n=0}^{\infty} E_n\right).$$

Or: $\mathbb{P}(\text{ultimate extinction}) = \mathbb{P}\left(\lim_{n \rightarrow \infty} E_n\right)$ i.e. $\mathbb{P}(\text{extinct by generation } \infty)$.

Note: By the Continuity Theorem (Chapter 2), and because $E_0 \subseteq E_1 \subseteq E_2 \subseteq \dots$, we have:

$$E_n = \{\text{extinct by generation } n\}.$$

$$\mathbb{P}(\text{ultimate extinction}) = \mathbb{P}\left(\lim_{n \rightarrow \infty} E_n\right) = \lim_{n \rightarrow \infty} \mathbb{P}(E_n).$$

Thus the probability of eventual extinction is the limit as $n \rightarrow \infty$ of the probability of extinction by generation n .

We will use the Greek letter Gamma (γ) for the probability of extinction: think of Gamma for 'all Gone'!

$$\gamma_n = \mathbb{P}(E_n) = \mathbb{P}(\text{extinct by generation } n)$$

$$\gamma = \mathbb{P}(\text{ultimate extinction}).$$

By the Note above, we have established that we are looking for:



$$\mathbb{P}(\text{ultimate extinction}) = \gamma = \lim_{n \rightarrow \infty} \gamma_n$$

$$\gamma$$

Extinction is Forever

Theorem 7.1: Let γ be the probability of ultimate extinction. Then

γ is the smallest non-negative solution of the equation $G(s) = s$, where $G(s)$ is the PGF of the family size distribution, Y . $\rightarrow \mathbb{E}(s^Y)$. i.e. $G(\gamma) = \gamma$.

To find the probability of ultimate extinction, we therefore:

- find the PGF of family size, Y : $G(s) = \mathbb{E}(s^Y)$.
- find values of s that satisfy $G(s) = s$;
- find the smallest of these values that is ≥ 0 . This is the required value, γ .

$G(\gamma) = \gamma$, and γ is the smallest value ≥ 0 for which this holds.

If $G(0) = 0$,
 $\gamma = 0$

$G(0) = \mathbb{P}(Y=0)$
 So $G(0) = 0 \Rightarrow \gamma = 0$,

Note: Recall that, for any (non-defective) random variable Y with PGF $G(s)$,

$$G(1) = \mathbb{E}(1^Y) = \sum_{y=0}^{\infty} 1^y \mathbb{P}(Y=y) = \sum_{y=0}^{\infty} \mathbb{P}(Y=y) = 1.$$

So $G(1) = 1$ always, and therefore there always exists a solution to $G(s) = s$ in the interval $[0, 1]$.

The required value γ is the smallest such solution ≥ 0 .

$(s-1)$ will always be a factor in solving $G(s) - s = 0$.

Before proving Theorem 7.1 we prove the following Lemma.

Lemma: Let $\gamma_n = \mathbb{P}(Z_n = 0)$. Then $\gamma_n = G(\gamma_{n-1})$.

Proof: If $G_n(s)$ is the PGF of Z_n ,

then $\mathbb{P}(Z_n = 0) = G_n(0)$. (True for any r.v. see Ch 4.)

So $\gamma_n = G_n(0)$.

Similarly, $\gamma_{n-1} = G_{n-1}(0)$.

Now $\gamma_n = G_n(0) = \underbrace{G(G(\dots G(0) \dots))}_{n \text{ times}} = G(\gamma_{n-1})$

$n \rightarrow \infty$
 $\gamma_n \rightarrow \gamma$
 $\gamma_{n-1} \rightarrow \gamma$
This says $\gamma = G(\gamma)$.

Proof of Theorem 7.1: We need to prove:

(i) $G(\gamma) = \gamma$.

(ii) γ is the smallest non-negative value for which $G(\gamma) = \gamma$.

That is, if $s \geq 0$ and $G(s) = s$, then $\gamma \leq s$.


$\therefore \gamma_n = G(\gamma_{n-1})$ as required. \square

Proof of (i):

From overleaf, $\gamma = \lim_{n \rightarrow \infty} \gamma_n = \lim_{n \rightarrow \infty} G(\gamma_{n-1})$ by Lemma

[because G is continuous] $= G(\lim_{n \rightarrow \infty} \gamma_{n-1})$
 $= G(\gamma)$ by def. of γ .

So $\gamma = G(\gamma)$, as required.



Proof of (ii):

Don't need to learn
→ might need to understand.

First note that $G(s)$ is an increasing function on $[0, 1]$:

$$G(s) = \mathbb{E}(s^Y) = \sum_{y=0}^{\infty} s^y \mathbb{P}(Y = y)$$

$$\Rightarrow G'(s) = \sum_{y=0}^{\infty} y s^{y-1} \mathbb{P}(Y = y)$$

sum of positive terms must be positive.

$$\Rightarrow G'(s) \geq 0 \text{ for } 0 \leq s \leq 1, \text{ so } G \text{ is increasing on } [0, 1].$$

$G(s)$ is increasing on $[0, 1]$ means that:

$$s_1 \leq s_2 \Rightarrow G(s_1) \leq G(s_2) \text{ for any } s_1, s_2 \in [0, 1].$$

The branching process begins with $Z_0 = 1$, so

$$\mathbb{P}(\text{extinct by generation } 0) = \gamma_0 = 0.$$

At any later generation, $\gamma_n = G(\gamma_{n-1})$ by Lemma.

Now suppose that $s \geq 0$ and $G(s) = s$. Then we have:

Set up an "imposter" solution, s .

by assumption $\leftarrow 0 \leq s \Rightarrow \gamma_0 \leq s$ (because $\gamma_0 = 0$)

$$\Rightarrow G(\gamma_0) \leq G(s) = s \text{ (by } \clubsuit \text{)}$$

i.e. $\gamma_1 \leq s$

$$\Rightarrow G(\gamma_1) \leq G(s) \text{ (by } \clubsuit \text{)}$$

i.e. $\gamma_2 \leq s$

\vdots

Thus $\gamma_n \leq s$ for all n .

So if $s \geq 0$ and $G(s) = s$, then $\gamma = \lim_{n \rightarrow \infty} \gamma_n \leq s$.



□

Example 1: Let $\{Z_0 = 1, Z_1, Z_2, \dots\}$ be a branching process with family size distribution $Y \sim \text{Binomial}(2, \frac{1}{4})$. Find the probability that the process will eventually die out.

Solution: Let $Q(s) = \mathbb{E}(s^Y)$. The probability of ultimate extinction is γ , where γ is the smallest solution ≥ 0 to the eqn.
 $Q(s) = s$. Need to solve $Q(s) = s$.

For $Y \sim \text{Bin}(n, p)$, the PGF is $Q(s) = (ps + q)^n$ (Attachment in Exam)
 So for $Y \sim \text{Bin}(2, \frac{1}{4})$, then $Q(s) = (\frac{1}{4}s + \frac{3}{4})^2$.

Need to solve $Q(s) = s$:

$$(\frac{1}{4}s + \frac{3}{4})^2 = s$$

$$(s + 3)^2 = 16s$$

$$s^2 - 10s + 9 = 0$$


$$(s-1)(s-9) = 0$$

$$\therefore s=1 \text{ or } s=9.$$

Trick! Know $s-1$ MUST BE A FACTOR!

γ is smallest such $s \geq 0$,

so $\gamma = 1$ $\therefore \mathbb{P}(\text{extinct}) = 1$. ☹️



Extinction is definite when the family size is $Y \sim \text{Bin}(2, \frac{1}{4})$.

$$2 * \frac{1}{4} = 0.5 = \mathbb{E}(\# \text{offspring}) \text{ per indiv.}$$



~~$E(Z_n) = \mu^n$~~ $\rightarrow \mu < 1$ then $E(Z_n) \rightarrow 0$ as $n \rightarrow \infty$.

Example 2: Let $\{Z_0 = 1, Z_1, Z_2, \dots\}$ be a branching process with family size distribution $Y \sim \text{Geometric}(\frac{1}{4})$. Find the probability that the process will eventually die out.

Solution: Let $G(s) = E(s^Y)$

$$= \frac{p}{1 - qs} \quad \text{for } Y \sim \text{Geo}(p)$$

$$= \frac{1/4}{1 - 3s/4} \quad \text{for } Y \sim \text{Geo}(\frac{1}{4})$$

$$\therefore G(s) = \frac{1}{4 - 3s}$$

Need to solve $G(s) = s$:

$$\Rightarrow \frac{1}{4 - 3s} = s$$

$$\Rightarrow 4s - 3s^2 = 1$$

$$3s^2 - 4s + 1 = 0$$

Trick! $(s-1)$ must be a factor!


$$\therefore (s-1)(3s-1) = 0$$

$$\Rightarrow s=1 \quad \text{or} \quad 3s-1=0 \Rightarrow s=\frac{1}{3}$$

γ = smallest such s that is ≥ 0 :

$$\text{So } \gamma = P(\text{extinction}) = \frac{1}{3}$$

Extinction is possible, but not definite,
When $Y \sim \text{Geo}(\frac{1}{4})$.



$$\hookrightarrow E(Y) = \frac{q}{p} = 3.$$

7.2 Conditions for ultimate extinction : i.e. when will $\gamma = 1$?

It turns out that the probability of extinction depends crucially on the value of μ , the mean of the family size distribution, Y . $\mu = \mathbb{E}Y$.
Some values of μ guarantee that the branching process will die out with probability 1. Other values guarantee that the probability of extinction will be strictly less than 1. We will see below that the threshold value is $\mu = 1$.

If the mean number of offspring per individual μ is more than 1 (so on average, individuals replace themselves plus a bit extra), then the branching process is not guaranteed to die out — although it might do. However, if the mean number of offspring per individual μ is 1 or less, the process is guaranteed to become extinct (unless $Y = 1$ with probability 1). The result is not too surprising for $\mu > 1$ or $\mu < 1$, but it is a little surprising that extinction is generally guaranteed if $\mu = 1$.

Theorem 7.2: Let $\{Z_0 = 1, Z_1, Z_2, \dots\}$ be a branching process with family size distribution Y . Let $\mu = \mathbb{E}(Y)$ be the mean family size distribution, and let γ be the probability of ultimate extinction. Then

- (i) If $\mu > 1$, then $\gamma < 1$: extinction is NOT guaranteed if $\mu > 1$.
- (ii) If $\mu < 1$, then $\gamma = 1$: extinction is GUARANTEED if $\mu < 1$.
- (iii) If $\mu = 1$, then $\gamma = 1$ UNLESS the family size is always constant at $Y = 1$. (very boring branching process)

Lemma: Let $G(s)$ be the PGF of family size Y . Then $G(s)$ and $G'(s)$ are strictly increasing for $0 < s < 1$, as long as Y can take values ≥ 2 .

Proof: $G(s) = \mathbb{E}(s^Y) = \sum_{y=0}^{\infty} s^y \mathbb{P}(Y = y)$.

So $G'(s) = \sum_{y=1}^{\infty} y s^{y-1} \mathbb{P}(Y = y) > 0$ for $0 < s < 1$,
 $2s^1 p_2 > 0$

because all terms are ≥ 0 and at least 1 term is > 0 (if $\mathbb{P}(Y \geq 2) > 0$).

Similarly, $G''(s) = \sum_{y=2}^{\infty} y(y-1) s^{y-2} \mathbb{P}(Y = y) > 0$ for $0 < s < 1$.
e.g. $2 \times 1 \times s^0 \times p_2 > 0$

So $G(s)$ and $G'(s)$ are strictly increasing for $0 < s < 1$. \square


as long as $\mathbb{P}(Y \geq 2) > 0$.

$G(s)$ is a banana!!



e.g. $p_2 > 2$

Note: When $G''(s) > 0$ for $0 < s < 1$, the function G is said to be **convex** on that interval.




$G''(s) > 0$ means that the gradient of G is constantly increasing for $0 < s < 1$.

Proof of Theorem 7.2: This is usually done graphically.

The graph of $G(s)$ satisfies the following conditions:

1. $G(s)$ is increasing and strictly convex (as long as Y can be ≥ 2)
2. $G(0) = P(Y=0) \geq 0$
3. $G(1) = 1$ (for any non-defective r.v.)
4. $G'(1) = E(Y) = \mu$. So the slope of the "banana" at 1 gives the crucial value μ .
5. The extinction prob, γ , is the smallest value ≥ 0 for which $G(s) = s$.




ie. banana crosses straight line.

Case (i): $\mu > 1$

When $\mu > 1$, the curve $G(s)$ is forced beneath the line $t=s$ at the point $s=1$.

The curve $G(s)$ has to cross the line $t=s$ again in order to meet the t -axis at $P(Y=0)$. So there MUST be a solution $\gamma < 1$ to the equation $G(s)=s$.




Case (ii): $\mu < 1$

When $\mu < 1$, the curve $G(s)$ is forced above the line $t=s$ for $s < 1$.

There is no possibility for the curve $G(s)$ to cross the line $t=s$ again before meeting the t -axis.

Thus there can be no solution < 1 to the equation $G(s)=s$, so $\gamma = 1$.

The exception is where Y can take only values 0 and 1, so $G(s)$ is not strictly convex (see Lemma). However, in that case $G(s) = p_0 + p_1 s$ is a straight line, giving the same result $\gamma = 1$.




Case (iii): $\mu = 1$

When $\mu = 1$, the situation is the same as for $\mu < 1$.

very boring The exception is where Y takes only the value 1. Then $G(s) = s$ for all $0 \leq s \leq 1$, so the smallest solution ≥ 0 is $\gamma = 0$.

Thus extinction is guaranteed for $\mu = 1$, **unless** $Y = 1$ with probability 1.



$$\mu < 1 \Rightarrow \gamma = 1.$$

Example 1: Let $\{Z_0 = 1, Z_1, Z_2, \dots\}$ be a branching process with family size distribution $Y \sim \text{Binomial}(2, \frac{1}{4})$, as in Section 7.1. Find the probability of eventual extinction.

Solution: $Y \sim \text{Bin}(2, \frac{1}{4}) \Rightarrow \mu = \mathbb{E}Y = 2 * \frac{1}{4} = \frac{1}{2} < 1.$
So by Thm 7.2, $\mu < 1 \Rightarrow \gamma = 1.$
 $\therefore \mathbb{P}(\text{ultimate extinction}) = 1.$

(Longer calculation in § 7.1 was not necessary.)

Example 2: Let $\{Z_0 = 1, Z_1, Z_2, \dots\}$ be a branching process with family size distribution $Y \sim \text{Geometric}(\frac{1}{4})$, as in Section 7.1. Find the probability of eventual extinction.

Solution: $Y \sim \text{Geo}(\frac{1}{4}) \Rightarrow \mu = \mathbb{E}Y = \frac{3/4}{1/4} = 3 > 1.$
So we know $\gamma < 1$,
but we still need the calculation in § 7.1 to find what γ is—
(Solve $G(s) = s \Rightarrow \gamma = \frac{1}{3}.$)

Note: The mean μ of the offspring distribution Y is known as the **criticality parameter**.

- If $\mu < 1$, extinction is definite ($\gamma = 1$). The process is called subcritical. Note that $\mathbb{E}(Z_n) = \mu^n \rightarrow 0$ as $n \rightarrow \infty$.
- If $\mu = 1$, extinction is definite unless $Y \equiv 1$. The process is called critical. Note that $\mathbb{E}(Z_n) = \mu^n = 1 \forall n$, even though extinction is definite.
 $\gamma = 1$ but $\mathbb{E}(\text{time to extinction}) = \infty$
- If $\mu > 1$, extinction is not definite ($\gamma < 1$). The process is called supercritical. Note that $\mathbb{E}(Z_n) = \mu^n \rightarrow \infty$ as $n \rightarrow \infty$.



BAD Luck ! your population is SCHEDULED to become Extinct!

But how long have you got...?

Thurs 3:30pm - 4:30pm
14th. PLT1

7.3 Time to Extinction

Suppose the population is doomed to extinction — or maybe it isn't. Either way, it is useful to know how long it will take for the population to become extinct. This is the distribution of T , the number of generations before extinction. For example, how long do we expect a disease epidemic like SARS to continue? How long have we got to organize ourselves to save the kakapo or the tuatara before they become extinct before our very eyes?



1. Extinction by time n generation.

The branching process is extinct by time n if $Z_n = 0$.

Thus the probability that the process has become extinct by time n is:

$$\mathbb{P}(Z_n = 0) = G_n(0) = \gamma_n.$$

Note: Recall that $G_n(s) = \mathbb{E}(s^{Z_n}) = \underbrace{G(G(G(\dots G(s)\dots)))}_{n \text{ times}}.$

There is no guarantee that the PGF $G_n(s)$ or the value $G_n(0)$ can be calculated easily. However, we can build up $G_n(0)$ in steps:

$$\begin{aligned} \text{e.g. } G_2(0) &= G(G(0)), & \text{or } G_4(0) &= G_2(G_2(0)) \\ \text{then } G_3(0) &= G(G_2(0)) & & \text{etc.} \end{aligned}$$

2. Extinction at time n

Let T be the exact time of extinction. That is, $T = n$ if generation n is the first generation with no individuals:

$$T = n \iff Z_n = 0 \text{ AND } Z_{n-1} > 0.$$


Now by the Partition Rule,

$$\mathbb{P}(Z_n = 0 \text{ AND } Z_{n-1} > 0) + \mathbb{P}(Z_n = 0 \text{ AND } Z_{n-1} = 0) = \mathbb{P}(Z_n = 0) \quad (*)$$

But the event $\{Z_n = 0 \cap Z_{n-1} = 0\}$ is the event that the process is extinct by generation $n - 1$ AND it is extinct by generation n . However, we know it will always be extinct by generation n if it is extinct by generation $n - 1$, so the $Z_n = 0$ part is redundant. So

$$\mathbb{P}(\underbrace{Z_n = 0}_{\text{redundant}} \cap Z_{n-1} = 0) = \mathbb{P}(Z_{n-1} = 0) = G_{n-1}(0)$$

Similarly, $\mathbb{P}(Z_n = 0) = G_n(0)$



So $(*)$ gives:

$$\mathbb{P}(T = n) = \mathbb{P}(Z_n = 0 \cap Z_{n-1} > 0) = G_n(0) - G_{n-1}(0)$$

$$\therefore \boxed{\mathbb{P}(T = n) = \gamma_n - \gamma_{n-1} = G_n(0) - G_{n-1}(0).}$$

This gives the distribution of T , the exact time at which extinction occurs.

Example: Binary splitting. Suppose that the family size distribution is

$$Y = \begin{cases} 0 & \text{with probability } q = 1 - p, \\ 1 & \text{with probability } p. \end{cases}$$

Find the distribution of the time to extinction.



Ass 5 \rightarrow directly into Exam.

Solution:

Consider

Ass 3 harder Qs.
PGF. and before.

$Y \sim \text{Bin}(1, p)$

$$\underline{G(s)} = \mathbb{E}(s^Y) = qs^0 + ps^1 = \underline{q + ps}.$$

$$\underline{G_2(s)} = G(G(s)) = q + p(q + ps) = \underline{q(1 + p) + p^2s}.$$

$$\underline{G_3(s)} = G(G_2(s)) = q + p(q + pq + p^2s) = \underline{q(1 + p + p^2) + p^3s}.$$

$$\vdots$$

$$\underline{G_n(s)} = q(1 + p + p^2 + \dots + p^{n-1}) + p^n s.$$

Thus time to extinction, T , satisfies

$$\mathbb{P}(T = n) = G_n(0) - G_{n-1}(0)$$

$$= q(1 + p + p^2 + \dots + p^{n-1}) - q(1 + p + p^2 + \dots + p^{n-2})$$

$$\underline{\mathbb{P}(T-1 = n-1)} = \underline{qp^{n-1}} \quad \text{for } n = 1, 2, \dots$$

$T \sim 1 + \text{Geo}(q).$

Thus

$$\underline{T - 1 \sim \text{Geometric}(q)}.$$

It follows that $\mathbb{E}(T - 1) = \frac{p}{q}$, so

$$\mathbb{E}(T) = 1 + \frac{p}{q} = \frac{1 - p + p}{q} = \frac{1}{q}.$$

Note: The expected time to extinction, $\mathbb{E}(T)$, is:

- finite if $\mu < 1$;
- infinite if $\mu = 1$ (despite extinction being definite), if σ^2 is finite;
- infinite if $\mu > 1$ (because with positive probability, extinction never happens).

(Results not proved here.)

For Reading/Interest.

(Higher priorities this semester).

7.4 Case Study: Geometric Branching Processes

Recall that $G_n(s) = \mathbb{E}(s^{Z_n}) = \underbrace{G\left(G\left(G\left(\dots G(s)\dots\right)\right)\right)}_{n \text{ times}}.$

In general, it is not possible to find a closed-form expression for $G_n(s)$. We achieved a closed-form $G_n(s)$ in the Binary Splitting example (page 134), but binary splitting only allows family size Y to be 0 or 1, which is a very restrictive model.

The only non-trivial family size distribution that allows us to find a closed-form expression for $G_n(s)$ is the

When family size $Y \sim \text{Geometric}(p)$, we can do the following:

- Derive a closed-form expression for $G_n(s)$, the PGF of Z_n .
- Find the probability distribution of the **exact time of extinction, T** : not just the probability that extinction will occur at some unspecified time (γ).
- Find the **full probability distribution of Z_n** : probabilities $\mathbb{P}(Z_n = 0)$, $\mathbb{P}(Z_n = 1)$, $\mathbb{P}(Z_n = 2)$, \dots .

With $Y \sim \text{Geometric}(p)$, we can therefore calculate just about every quantity we might be interested in for the branching process.

1. Closed form expression for $G_n(s)$

Theorem 7.4: Let $\{Z_0 = 1, Z_1, Z_2, \dots\}$ be a branching process with family size distribution $Y \sim \text{Geometric}(p)$. The PGF of Z_n is given by:

$$G_n(s) = \mathbb{E}(s^{Z_n}) = \begin{cases} \frac{n - (n-1)s}{n+1 - ns} & \text{if } p = q = 0.5, \\ \frac{(\mu^n - 1) - \mu(\mu^{n-1} - 1)s}{(\mu^{n+1} - 1) - \mu(\mu^n - 1)s} & \text{if } p \neq q, \text{ where } \mu = \frac{q}{p}. \end{cases}$$

Proof (sketch):

The proof for both $p = q$ and $p \neq q$ proceed by mathematical induction. We will give a sketch of the proof when $p = q = 0.5$. The proof for $p \neq q$ works in the same way but is trickier.

Consider $p = q = \frac{1}{2}$. Then

$$G(s) = \frac{p}{1 - qs} = \frac{\frac{1}{2}}{1 - \frac{s}{2}} = \frac{1}{2 - s}.$$

Using the Branching Process Recursion Formula (Chapter 6),

$$G_2(s) = G(G(s)) = \frac{1}{2 - G(s)} = \frac{1}{2 - \frac{1}{2-s}} = \frac{2-s}{2(2-s) - 1} = \frac{2-s}{3-2s}.$$

The inductive hypothesis is that $G_n(s) = \frac{n - (n-1)s}{n+1 - ns}$, and it holds for $n = 1$ and $n = 2$. Suppose it holds for n . Then

$$\begin{aligned} G_{n+1}(s) &= G_n(G(s)) = \frac{n - (n-1)G(s)}{n+1 - nG(s)} = \frac{n - (n-1)\left(\frac{1}{2-s}\right)}{n+1 - n\left(\frac{1}{2-s}\right)} \\ &= \frac{(2-s)n - (n-1)}{(2-s)(n+1) - n} \\ &= \frac{n+1 - ns}{n+2 - (n+1)s}. \end{aligned}$$

Therefore, if the hypothesis holds for n , it also holds for $n+1$. Thus the hypothesis is proved for all n . \square

2. Exact time of extinction, T

Let $Y \sim \text{Geometric}(p)$, and let T be the exact generation of extinction.

From Section 7.3,

$$\mathbb{P}(T = n) = \mathbb{P}(Z_n = 0) - \mathbb{P}(Z_{n-1} = 0) = G_n(0) - G_{n-1}(0).$$

By using the closed-form expressions overleaf for $G_n(0)$ and $G_{n-1}(0)$, we can find $\mathbb{P}(T = n)$ for any n .

3. Whole distribution of Z_n

From Chapter 4, $\mathbb{P}(Z_n = r) = \frac{1}{r!} G_n^{(r)}(0)$.

Now our closed-form expression for $G_n(s)$ has the same format regardless of whether $\mu = 1$ ($p = 0.5$), or $\mu \neq 1$ ($p \neq 0.5$):

$$G_n(s) = \frac{A - Bs}{C - Ds}.$$

(For example, when $\mu = 1$, we have $A = D = n$, $B = n - 1$, $C = n + 1$.) Thus:

$$\mathbb{P}(Z_n = 0) = G_n(0) = \frac{A}{C}$$

$$G_n'(s) = \frac{(C - Ds)(-B) + (A - Bs)D}{(C - Ds)^2} = \frac{AD - BC}{(C - Ds)^2}$$

$$\Rightarrow \mathbb{P}(Z_n = 1) = \frac{1}{1!} G_n'(0) = \frac{AD - BC}{C^2}$$

$$G_n''(s) = \frac{(-2)(-D)(AD - BC)}{(C - Ds)^3} = \frac{2D(AD - BC)}{(C - Ds)^3}$$

$$\Rightarrow \mathbb{P}(Z_n = 2) = \frac{1}{2!} G_n''(0) = \left(\frac{AD - BC}{CD} \right) \left(\frac{D}{C} \right)^2$$

\vdots

$$\Rightarrow \mathbb{P}(Z_n = r) = \frac{1}{r!} G_n^{(r)}(0) = \left(\frac{AD - BC}{CD} \right) \left(\frac{D}{C} \right)^r \quad \text{for } r = 1, 2, \dots$$

(Exercise)

This is very simple and powerful: we can substitute the values of A, B, C , and D to find $\mathbb{P}(Z_n = r)$ or $\mathbb{P}(Z_n \leq r)$ for any r and n .

Note: A Java applet that simulates branching processes can be found at:

http://www.dartmouth.edu/~chance/teaching_aids/books_articles/probability_book/bookapplets/chapter10/Branch/Branch.html